

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/284025185>

Estimating DSSAT Cropping System Cultivar-Specific Parameters Using Bayesian Techniques

Article · January 2011

DOI: 10.2134/advagricsystmodel2.c13

CITATIONS

103

READS

1,271

6 authors, including:



James W Jones

University of Florida

470 PUBLICATIONS 30,659 CITATIONS

[SEE PROFILE](#)



Jianqiang He

Northwest A & F University

102 PUBLICATIONS 2,044 CITATIONS

[SEE PROFILE](#)



Cheryl H Porter

University of Florida

80 PUBLICATIONS 8,602 CITATIONS

[SEE PROFILE](#)

Estimating DSSAT Cropping System Cultivar-Specific Parameters Using Bayesian Techniques

James W. Jones
Jianqiang He
Kenneth J. Boote
Paul Wilkens
C.H. Porter
Z. Hu




Abstract

Crop models are highly useful for simulating crop and soil processes in response to variations in climate and management. However, if one wishes to simulate a crop's performance in a specific soil and climate for a particular set of management inputs, cultivar-specific parameters (CSPs) are needed because of the genetic variations among cultivars of any crop. In this chapter, we summarized methods that have been used to estimate CSPs for the CERES and CROPGRO-based models in the Decision Support System for Agrotechnology Transfer (DSSAT) cropping system model. We primarily described a Bayesian parameter estimation procedure (the Generalized Likelihood Uncertainty Estimation, or GLUE) for use in estimating CSPs in DSSAT. The procedure is simple to use, requiring only that users select a crop, a cultivar, and the data for use in the estimation procedure from a list of data available for that cultivar in the DSSAT system. Results are displayed for users to view and copy to the standard cultivar file in DSSAT for the crop involved. The procedure does require a large number of model runs; we recommend 6000 but users can optionally change this number. Two cultivars, 'Prisma' maize (*Zea mays* L.) and 'Williams' soybean [*Glycine max* (L.) Merr.] were selected to demonstrate the performance of DSSAT GLUE program. For Prisma maize, two experiments conducted in Zaragoza, Spain in 1995 and 1996 were selected; for Williams soybean, three experiments individually conducted in Wooster, OH and Gainesville, FL, were selected for the demonstration. Results showed that the GLUE method performed better than the arbitrary default CSPs and slightly better than the hand-calibrated CSPs in simulating these maize and soybean cultivars when using one time measurements, such as phenology dates, final dry matter yield, maximum leaf area index, and grain yield. For example, in the Prisma maize experiments in Zaragoza, Spain in 1995 and 1996, the average relative absolute error (RAE) values between the simulated and measured output variables were only 3 and 8%, respectively, while they were between 4 and 10% for hand-calibrated CSPs and above 16% for the default CSPs.

J.W. Jones (jimj@ufl.edu), C.H. Porter, and Z. Hu, Dep. of Agricultural and Biological Engineering; K.J. Boote, Agronomy Dep., Univ. of Florida, Gainesville, FL 32611. Jianqiang He, INRA, UMR1095, Génétique, Diversité et Ecophysiologie des Céréales, F-63100 Clermont-Ferrand, France; P. Wilkens, International Fertilizer Development Center, Muscle Shoals, AL 35662.

doi:10.2134/advagriscystmodel2.c13

Copyright © 2011, ASA, CSSA, SSSA, 5585 Guilford Rd., Madison, WI 53711-5801, USA.
Methods of introducing system models into agricultural research. L.R. Ahuja and L. Ma (ed.)
Advances in Agricultural Systems Modeling Series 2.



Crop models are highly useful for simulating crop and soil processes in response to variations in climate and management. The basic concept is that simulating crop growth and yield using dynamic crop models will produce results that represent how a real crop growing under specific environmental and management conditions would perform. Furthermore, if simulated results are sufficiently accurate, one can perform experiments using the models to evaluate likely responses to climate, soil properties, and crop and soil management. However, there are practical limitations that must be considered before making use of this approach in any study. One main limitation is that crop models do not contain all of the factors in the field that may influence crop yield. For example, crop diseases, weeds, and spatial variability of soils and management implementation can cause large differences in yield, and these factors are seldom included in crop simulation analyses. Another limitation is that inputs must be accurate or else simulated outputs are unlikely to match observations from the field. Attempts to evaluate the predictability of a crop model thus require that weather, management, and soil inputs are measured in the field where the evaluation experiments are conducted. Furthermore, model evaluation experiments would ideally be designed to eliminate yield-reducing factors that are not included in the model. Finally, parameters that are used to model the dynamics of soil and crop processes need to be accurate for comparison with observed field data. For example, if one uses a crop model to simulate crop yield responses to water or N management using incorrect soil water parameters, results will show that the model fails to mimic results from field experiments or, more problematically, will provide results that may mislead researchers or other model users.

If, however, one wishes to simulate crop growth and yield for large areas in which soils and climate vary, the input soil, weather, and management conditions should represent the spatial variability that exists over the area to provide reliable estimates at aggregate scales. In this case, many possible fields may be simulated without an attempt to mimic any particular real field. The objectives of such studies may be to compare aggregate effects of crops, varieties, and management systems under the range of soil and climate conditions that existing in an area. Options can be evaluated for changing crop management in an area to achieve a goal, such as to study management options for adapting to potential

changes in climate. For these broad studies, model inputs need to represent the spatial variability that exists over the landscape, and crop and variety parameters should accurately represent those of a particular field from which observations were collected for comparisons with the model.

In this chapter, we focus on methods for estimating parameters for the DSSAT cropping system model (Tsuji et al., 1994; Jones et al., 2003; Hoogenboom et al., 2003; Boote et al., 2003) using data collected from field studies on real cropping systems. When field observations are used to estimate model parameters, however, the resulting parameters may not be transferrable to other conditions. This model transferability depends on its robustness, the parameters used in model equations, and the quality of observations used as model inputs and response variables. Parameters in models are often highly related to their testing conditions and are less universal than expected. Therefore, all crop models should be calibrated and validated for the environment of interest if results are to be credible (Timsina and Humphreys, 2006).

Emphasis is given on estimating the so-called genetic coefficients (Ritchie et al., 1986; Hunt et al., 1993; Boote et al., 2003), or what are more accurately referred to as cultivar-specific parameters (CSPs). The implicit assumption in the models is that there are parameters for a given crop species that remain the same for all cultivars and that there are parameters that vary among cultivars and allow simulating differences in yield or other traits when different cultivars are grown in the same environments and management conditions (Bertin et al., 2009; Boote et al., 2001). This is important because there are many existing cultivars, and new ones are released each year. Because the CSPs are not known for most cultivars, model users need to estimate them using field data. The emphasis in this chapter on CSPs should not be interpreted as a statement that they are more important than soil parameters. As stated above, when one attempts to evaluate simulated results with field observations, soil physical and chemical properties and initial conditions are very important. In some cases, crop yield responses are more strongly affected by uncertainties in soil parameters than by CSPs (Jones et al., 2010; Mavromatis et al., 2001).

One should not expect a crop model to simulate soil water and nutrient dynamics or growth and yield of a crop accurately unless soil physical and chemical properties and CSPs are accurate. The methods presented in this chapter can also be used to estimate soil parameters if appropriate soil measurements are taken in the field where an experiment is conducted (He et al., 2009), but the procedure implemented in DSSAT only estimates CSPs. Furthermore, a model may not be sensitive to some of its parameters, and in such cases, one may need to put less effort into estimating those parameters than others. A sensitivity analysis is

generally recommended before estimating parameters for models (e.g., see Jawitz et al., 2008; Monod et al., 2006; Makowski et al., 2006; Muñoz-Carpena et al., 2007, 2010; Saltelli et al., 2004). However, we used prior experience to select the CSPs to estimate in the DSSAT implementation, so it is not necessary for users to do this before using the DSSAT GLUE program.

The objectives of this chapter are to summarize methods that have been used to estimate CSPs for the CERES and CROPGRO-based models in DSSAT and to present a Bayesian parameter estimation procedure (the Generalized Likelihood Uncertainty Estimation, or GLUE) that is in the latest release of DSSAT (version 4.5, Hoogenboom et al., 2010). Some results are presented to highlight the capabilities and limitations of this approach.

General Parameter Estimation Approaches

Model parameter estimation, sometimes referred to as calibration, is the process of estimating parameters to obtain a match between observed and simulated system behavior (Oliva, 2003). This requires a set of observed data from the real system, the model that simulates the system's behavior, a criterion for determining the best parameters, and a method to determine the best parameter set. Generally, the criterion is to minimize the error between observed and simulated variables. Many methods have been developed and used, such as conjugate gradient-descent search (Bhalla and Bower, 1993), stochastic-search (Foster et al., 1993), genetic algorithms (Baldi et al., 1998; Eichler-West and Wilcox, 1997; Vanier and Bower, 1996), and simulated annealing (Baldi et al., 1998; and Vanier and Bower, 1996). Perhaps the most common approach is trial and error (Wallach et al., 2001). Various parameter values are tested until a set of values is found that gives an acceptable fit to the data. For example, when Müller et al. (2003) calibrated the DAISY model to simulate decomposition of plant residues in soil, they used a stepwise trial and error process. The criterion of each iteration step was either to minimize the root mean square error (RMSE) or to maximize modeling efficiency (EF). The trial and error process has been criticized because it is unreliable and difficult to replicate (Lyneis and Pugh, 1996). Manual calibration may also be very tedious and time-consuming, depending on the number of model parameters and the degree of parameter interaction. Thus, a great deal of research has been directed to development of more effective and efficient automatic calibration procedures (Madsen et al., 2002).

There are two general approaches used to estimate parameters, frequentist and Bayesian (Makowski et al., 2006). The frequentist approach uses estimation methods to approximate assumedly true and fixed parameter values by using a

sample of data. Prior information on parameter values is not taken into account. The use of a frequentist method gives a single, deterministic estimate of each parameter. In contrast, Bayesian methods estimate parameters from two types of information, a sample of data (like the frequentist method) and prior information about parameter values. The results of Bayesian methods are probability distributions of parameter values. All Bayesian methods proceed in two steps. The first step is to define a “prior” parameter probability distribution based on literature or expert knowledge. The second step involves calculating a new parameter probability distribution from both the prior distribution and the available data set. This new distribution, termed the *posterior distribution*, is computed by using Bayes theorem (Makowski et al., 2006).

Bayesian methods are becoming increasingly popular for estimating parameters for complex mathematical models (e.g., Campbell et al., 1999) because these methods provide a coherent framework for dealing with uncertainty. This is also due to the increase in speed of computers and the development of new algorithms (Malakoff, 1999). One commonly used Bayesian method is the GLUE method (Beven and Binley, 1992; Franks et al., 1998; Shulz et al., 1999). The GLUE method assumes that, in the case of large models with many parameters, there is no exact inverse solution. Hence, the estimation of a unique set of parameters, which optimizes a goodness-of-fit criterion given the observations, is not possible (Romanowicz and Beven, 2006). The main principle of this method is to discretize the parameter space by generating a large number of parameter values from the prior distribution. Likelihood values are then calculated for each parameter set using field observations. Probabilities, an empirical posterior distribution of the parameters, are calculated using Bayes’ equation.

Since it was introduced in 1992, the GLUE framework has found widespread application in environmental modeling (Blasone et al., 2008). The popularity of GLUE is largely due to its conceptual simplicity, relative ease of implementation and use, and its ability to handle different error structures and models without major modifications to the method itself (Blasone et al., 2008). Example applications include those for rainfall–runoff (Beven and Binley, 1992; Freer et al., 1996; Lamb et al., 1998), soil erosion (Brazier et al., 2001), tracer dispersion in a river reach (Hankin et al., 2001), groundwater and well capture zone delineation (Feyen et al., 2001; Jensen, 2003), unsaturated zone (Mertens et al., 2004), flood inundation (Romanowicz et al., 1996; Aronica et al., 2002), land–surface–atmosphere interactions (Franks et al., 1997), soil freezing and thawing (Hansson and Lundin, 2006), crop yields and soil organic carbon (Wang et al., 2005), ground radar rainfall estimation (Tadesse and Anagnostou, 2005), and distributed hydrology (McMichael et al., 2006; Muleta and Nicklow, 2005). He et

al. (2009, 2008) used the GLUE method to estimate soil parameters and CSPs in the DSSAT CERES-Maize model.

There are a number of other Bayesian methods for estimating parameters, including the Metropolis–Hastings method (Makowski et al., 2002). We implemented the Metropolis–Hastings method in one study and found that it is more difficult to implement, but more efficient than GLUE (Hu and Jones, unpublished data, 2010). The major difference between these methods is that the Metropolis–Hastings method uses a Markov Chain–Monte Carlo (MCMC) parameter search method, whereas the GLUE method uses a Monte Carlo random search method. In this chapter, we present the GLUE method that is implemented in the latest release of the DSSAT software (version 4.5, Hoogenboom et al., 2010).

Materials and Methods

Overview of DSSAT

The DSSAT is a software system that combines crop models with observed data from field trials and tools that help users enter data from experiments, evaluate the models, estimate CSPs, conduct sensitivity analyses, analyze economic risk and uncertainty of alternative management options, and graphically present simulated and observed results (Uehara, 1998; Jones et al., 1998; Hoogenboom et al., 2003). One of the unique features of this system is that it has databases that connect to the crop models and are used for evaluating model performance and estimating CSPs. At the heart of DSSAT is the cropping system model (DSSAT-CSM), which incorporates all crops as modules using a single soil model (Jones et al., 2003). The DSSAT version 4.5 can simulate more than 20 crops, including maize, wheat (*Triticum aestivum* L.), rice (*Oryza sativa* L.), sorghum [*Sorghum bicolor* (L.) Moench], soybean, and peanut (*Arachis hypogaea* L.). The CSM simulates growth, development, and yield of a crop growing on a uniform area of land under specified management. The dynamics of soil water, carbon, nitrogen and phosphorus that take place in the cropping system through time are also simulated. The model is structured using the modular approach described by Jones et al. (2001) and Porter et al. (2000) and consists of a main driver program, a land unit module, and primary modules for weather, management, soil, plant, and soil–plant–atmosphere interface components.

A number of methods have been used in the past to estimate CSPs for the different crops in the DSSAT-CSM. In many cases, trial and error methods are used to select a set of parameters that visually fits the observed data and/or produces an acceptably low RMSE between simulated and observed variables, often using a combination of time-series and end-of-season data concurrently (e.g., Boote,

1999; Boote et al., 2003). Other methods include the Simplex method (Grimm et al., 1993), Simulated Annealing (Mavromatis et al., 2001, 2002), and the K-Nearest Neighbor approach (Bannayan and Hoogenboom et al., 2008). A CSP estimation tool was developed by Hunt et al. (GENCALC, Hunt et al., 1993) and integrated into DSSAT v3.0, and v3.5. This tool automated a systematic search of parameters that minimized RMSE between simulated and observed variables; the criteria for selecting parameter values depended on the parameter being estimated in a sequential search process. A new updated version of this tool is also available in DSSAT v4.5.

Implementation of GLUE in DSSAT

The GLUE CSP estimation method was integrated into DSSAT using the R language (R Development Core Team, 2009; <http://www.R-project.org>, verified 13 May 2011), a free software environment for statistical computing and graphics. The program is simple to use in that users only have to select a crop, a cultivar (from a list of cultivars included in the DSSAT database for that crop), and the treatments from the various experiments in which that cultivar was grown (Fig. 13–1). The ranges of CSPs are stored in a file so that users do not have to specify these values. Optionally, users can choose to estimate only phenology CSPs, only growth CSPs, or both of these simultaneously, and set the number of runs to make. The interface also allows users to view the final estimated parameters and their distributions. The GLUE calculations are made using the GLUE R program, which calls the DSSAT-CSM, connects the data to the model, generates all parameter sets, and performs all of the necessary calculations to select the CSPs.

Theoretical Basis of the GLUE Method in DSSAT v4.5

Although users do not need to know all of the calculations that are done to arrive at the estimated CSPs when they use the GLUE program, we summarize the main theoretical basis for the method and the steps used internally in the program. In the GLUE procedure, a likelihood function is used as the criterion for estimating CSPs similar to that used by He et al. (2009). A likelihood value is computed for each observation, given each particular set of randomly generated CSP sets. The likelihood function is the product of these individual likelihood values. In DSSAT, the Gaussian likelihood function (Eq. [1]) (Beven and Binley, 1992; Romanowicz et al., 1994, 1996; Makowski et al., 2002; He et al. (2009, 2008, 2010) was used:

$$L[\theta_i | O] = \prod_{j=1}^M \frac{1}{\sqrt{2\pi\sigma_o^2}} \exp\left\{-\frac{[O_j - Y(\theta_i)]^2}{2\sigma_o^2}\right\}, \quad (i = 1, 2, 3 \dots N) \quad [1]$$




Fig. 13–1. Screen shots of the DSSAT GLUE program user interface. (Top) Users first select a crop, then a cultivar, (middle) then select which treatments they wish to use for estimating the cultivar-specific parameters (CSPs), and (bottom) then run the procedure. After the program finishes, users view the coefficients and can copy and paste the estimated CSPs into the crop cultivar file. Users may also view the standard deviations of CSP estimates from this last screen.

where θ_i is the i th parameter set, N is the total number of parameter sets generated by the program, $Y(\theta_i)$ is the model output using parameter set θ_i , O is the observation; O_j is the j th observation of O ; σ_o^2 is the variance of model error, and M is the number of observations. The probability p_i of each parameter set is computed with the equation (He et al., 2009):

$$p(\theta_i) = \frac{L(\theta_i | Y)}{\sum_{i=1}^N L(\theta_i | Y)} \quad [2]$$

where $p(\theta_i)$ is the probability or likelihood weight of the i th parameter set θ_i , and $L(\theta_i | Y)$ is the likelihood value of parameter set θ_i , given observations Y .

The GLUE methodology (Beven and Binley, 1992) thus develops an approximate discrete posterior probability distribution, designated by (θ_i, p_i) , $i = 1, \dots, N$, $\sum_{i=1}^N p_i = 1$, where p_i is the probability associated with the parameter vector θ_i , and N is the total number of generated parameter vectors.

Equations [1] and [2] are used to construct the posterior distributions of the CSPs. The implementation in DSSAT uses two iterations of GLUE, one to estimate phenological development parameters and the others to estimate growth parameters. This was done for practical reasons, mainly due to the time required when estimating all parameters simultaneously. This can be done in the DSSAT models because development is largely independent of growth but not vice versa. Thus, the development parameters are first estimated followed by those that affect biomass growth and yield in a two-step sequence. Thus, there are two posterior distributions, one for each step.

The main steps of the GLUE procedure in DSSAT are based on Beven and Binley (1992) and are summarized as follows:

1. *Develop prior parameter distributions.* In this study, the CSPs of two crops (maize and soybean) in the DSSAT database (Hoogenboom et al., 2003) were analyzed to determine the prior range of each parameter. Because we do not have additional information, the prior distributions are assumed to be independent and uniformly distributed between the minimum and maximum values across all cultivars previously calibrated for each crop.
2. *Generate random parameter sets from the prior parameter distributions.* A large number (e.g., 6000) of parameter sets are created by randomly generating each CSP in each of 6000 CSP vectors independently, according to the prior uniform distribution of each CSP. The number of runs can be modified by the user, but at least 3000 runs are recommended (He, 2008) to ensure that CSPs are each estimated accurately and the calculated posterior distributions are reliable.

3. *Run the model with the randomly generated parameter sets.* The model is run with the parameter sets generated above. The standard genetic input file is changed to simulate every random parameter set in sequence. Model outputs (e.g., anthesis date, maturity date, dry matter yield, leaf area index, and leaf number, which could be selected by model users) for each parameter set are tabulated for use in the GLUE likelihood calculations.
4. *Calculate the likelihood values.* The observations (Y) from the selected data provided by model users are used along with the corresponding simulated outputs to compute the likelihood value, $L(\theta_i|Y)$, for each generated parameter vector θ_i .
5. *Construct posterior distribution and statistics.* The pairs of parameter sets and probabilities, (θ_i, p_i) , $i = 1, \dots, N$, are computed and used to construct the posterior distribution and to compute the mean and variance of the selected parameters using following equations:

$$\hat{\mu}_{\text{post}}(\theta) = \sum_{i=1}^N p(\theta_i) \cdot \theta_i \quad [3]$$

$$\hat{\sigma}_{\text{post}}^2(\theta) = \sum_{i=1}^N p(\theta_i) \cdot (\theta_i - \hat{\mu}_{\text{post}})^2 \quad [4]$$

where $\hat{\mu}_{\text{post}}(\theta)$ and $\hat{\sigma}_{\text{post}}^2(\theta)$ are the estimated mean and variance of the posterior distribution of parameters θ , $p(\theta_i)$ is the probability of the i th parameter set θ_i calculated by Eq. [1], and N is the number of random parameter sets.

Application of GLUE to Estimate Cultivar-Specific Parameters of Maize and Soybean

Cultivar-specific parameters were estimated for maize and soybean to demonstrate the method's performance and characteristics of results that were obtained. In particular, we show observed and simulated values derived from the CSPs selected using the GLUE method. We also present uncertainties in the estimated CSPs and in simulated outputs.

Table 13-1 lists the CSPs of the CERES-Maize model in DSSAT (Jones et al., 2003) that are estimated in the GLUE procedure. The CSPs P1, P2, and P5 determine the timing of phenological events, such as anthesis date and maturity date of maize. Coefficients G2 and G3 control the yield-related outputs, such as grain dry matter yield, grain size, and canopy weight. Our two-step procedure requires that all CSPs estimated in the first step be completely determined by phenological development data. The CSPs estimated in the first step plus all other CSPs in the second step affect growth responses. Phenological development is assumed to be independent from growth in the model, but growth is affected by phenologi-

Table 13–1. Cultivar-specific parameters in the DSSAT CERES-Maize model (Jones et al., 2003) that are estimated in the DSSAT GLUE procedure.

Coefficient	Minimum	Maximum	GLUE flag [†]	Definition
P1	140	365	1	Degree days (base 8°C) from emergence to end of juvenile phase
P2	0.0	1.0	1	Photoperiod sensitivity coefficient (0–1.0)
P5	600	990	1	Degree days (base 8°C) from silking to physiological maturity
G2	500	908	2	Potential kernel number
G3	5	15	2	Potential kernel growth rate mg/(kernel d)

[†] GLUE flag is an indicator to show in which round of the procedure the parameter will be estimated.

cal development. This assumption is reasonable unless major stresses occur. The CERES-Maize CSP PHINT is not included in the procedure because its value is similar across many cultivars and because it influences both phenological development and yield. PHINT was assumed to be 48.0 for all cultivars.

Table 13–2 shows the CSPs for the CROPGRO-Soybean model in the DSSAT-CSM. There are 18 total CSPs in soybean (Boote et al., 2003), but the 11 coefficients in the table below are those that vary most among cultivars and determine differences in soybean cultivars' responses to their environments. The model is highly sensitive to these CSPs and less sensitive to the other ones when they are near their nominal values in the cultivar file available in DSSAT.

In this chapter, two cultivars (Prisma and Williams shown in Table 13–3) of two crops (maize and soybean, respectively) were selected as examples to evaluate the performance of DSSAT GLUE program. For Prisma maize, the experiments conducted in Zaragoza, Spain in 1995 and 1996 were used. There are a total of eight and nine treatments based on different irrigation levels in 1995 and 1996, respectively, but only the fully irrigated treatments were selected so that development and growth were influenced only by weather, not by water or nutrient stresses. Similarly, three experiments, conducted individually in Wooster, OH and Gainesville, FL, were selected for Williams soybean. Only irrigated treatments were used to estimate the CSPs. A summary of experimental treatment characteristics, including planting date, N application, irrigation, and available observations is presented in Table 13–3.

The monthly average weather data were summarized for growing season months for these five different experiments (Table 13–4). Generally, solar radiation was higher and rainfall (mm d⁻¹) was lower for this maize experiment site than the other sites. The Gainesville, FL, 1979, experiment had relatively higher minimum and maximum temperatures and lower rainfall, which is due the subtropical climate of Florida. In the Ohio soybean experiments, solar radia-

Table 13–2. Cultivar-specific parameters for the soybean crop in the DSSAT-CSM (from Boote et al., 2003) that are estimated in the GLUE procedure.

Coefficient	Minimum	Maximum	GLUE flag†	Definition
CSDL	11.78	14.6	1	Critical Short Day Length below which reproductive development progresses with no day length effect (for short day plants) (h)
PPSEN	0.129	0.349	1	Slope of the relative response of development to photoperiod with time (positive for short day plants) (1/h)
EM-FL	15.5	23.5	1	Time between plant emergence and flower appearance (R1) (PTD‡)
FL-SD	12	16	1	Time between first flower and first seed (R5) (PTD)
SD-PM	29.5	37.5	1	Time between first seed (R5) and physiological maturity (R7) (PTD)
LFMAX	1	1.4	2	Maximum leaf photosynthesis rate (at 30C, 350 ppm CO ₂ , high light, mg CO ₂ m ⁻² s ⁻¹)
SLAVR	350	425	2	Specific leaf area of cultivar under standard growth conditions (cm ² g ⁻¹)
SIZELF	140	230	2	Maximum size of full leaf (three leaflets) (cm ²)
WTPSD	0.155	0.195	2	Maximum weight per seed (g)
SFDUR	17	25.5	2	Seed filling duration for pod cohort at standard growth conditions (PTD)
SDPDV	1.7	2.44	2	Average seed per pod under standard growing conditions (no. pod ⁻¹)

† GLUE flag is an indicator to show in which round of the procedure the parameter will be estimated.

‡ PTD, photothermal days, comparable to calendar days if at optimum temperature and CSDL.

tion was higher and rainfall less than in Florida. Because irrigation was used in all of the experiments, water stress likely did not reduce yields relative to the potential yields.

The CSPs estimated using the GLUE procedure were compared with those from two other sources—an arbitrary default set of CSPs used to initialize the GLUE procedure and the hand-calibrated CSPs that were in the DSSAT v4.5 cultivar database. Field observations from the selected experiments (Table 13–3) in DSSAT were compared with the model-simulated output variables derived from the three sources of CSPs. The relative absolute error (RAE percentage; Eq. [5]) was used as a measure to evaluate the differences between simulated and observed variables for each set of CSPs:

$$\text{RAE} = \frac{|\hat{Y} - Y|}{Y} 100\% \quad [5]$$

where \hat{Y} and Y are simulated and measured variables, respectively.

Table 13–3. Summary of experiment details for the example maize and soybean cultivars and experiments in DSSAT v4.5 used for estimating Cultivar-Specific Parameters in this study (Hoogenboom et al. 2010). Also shown are the field observations used in each experiment.

Crop	Cultivar	Experiment	Experimental items	Details†	
Maize	Prisma	Zaragosa, Spain 1995	Treatment	Fully irrigated	
			Location	Zaragosa, Spain	
			Planting date	17 May 1995	
			N fertilizer	100 and 200 kg urea-N ha ⁻¹ 16 May 1995 and 16 June 1996	
			Irrigation	A sum of 568 mm water in 9 events	
			Available observations	ADAP, MDAP, HWAM, HWUM, H#UM, CWAM, LAIX, HIAM	
			Observations	CWAM, LAIX	
		Zaragosa, Spain 1996	Treatment	Fully irrigated	
			Location	Zaragosa, Spain	
			Planting date	16 May 1996	
			N fertilizer	100 and 200 kg urea-N ha ⁻¹ 15 May 1996 and 15 May 1996	
			Irrigation	A sum of 505 mm water in 8 events	
			Available observations	ADAP, MDAP, HWAM, HWUM, H#UM, CWAM, LAIX, HIAM	
			Observations	CWAM, LAIX	
Soybean	Williams	Wooster, OH 1988	Planting date	1 May 1988	
			N fertilizer	No N applied	
			Irrigation	A sum of 595 mm water in 18 events	
			Available observations	ADAP, PD1P, MDAP, HWAM, HWUM, CWAM, LAIX	
			Wooster, OH 1990	Planting date	30 Apr. 1990
				N fertilizer	No N applied
				Irrigation	Not irrigated (not necessary)
		Gainesville, FL 1979	Available	ADAP, PD1P, MDAP, HWAM, PWAM, PWAM, PWAM, H#AM, HWUM, H#UM, CWAM, BWAM, LAIX, HIAM, THAMS	
			Observations	HWUM, CWAM, LAIX	
			Planting date	15 Mar. 1979	
			N fertilizer	No N applied	
			Irrigation	A sum of 144 mm water in 14 events	
			Available observations	ADAP, PD1P, MDAP, HWAM, HWUM, CWAM, LAIX	

† These abbreviations represent the model output variables in DSSAT: ADAP, anthesis date (days after planting); MDAP, maturity date (days after planting); HWAM, yield at harvest maturity (kg [dm]/ha); HWUM, unit weight at maturity (g [dm]/unit); CWAM, tops weight at maturity (kg [dm]/ha); LAIX, maximum leaf area index; PD1P, first pod day (days after planting); PWAM, pod/ear/panicle weight at maturity (kg [dm]/ha).

Table 13–4. Monthly average weather data of the five experiments of maize and soybean (Hoogenboom et al. 2010).†

	Month	Apr.	May	June	July	Aug.	Sept.	Oct.
Zaragoza, 1995	Tmin	5.1	10.9	14.0	17.2	16.7	11.7	9.4
	Tmax	21.1	24.8	27.4	32.8	30.0	24.0	24.2
	Rain	2.0	1.2	0.3	0.1	0.5	0.3	0.1
	SRAD	22.1	21.5	25.1	25.6	22.9	17.8	12.5
Zaragoza, 1996	Tmin	6.4	10.0	14.3	15.9	16.0	11.8	8.1
	Tmax	19.3	23.2	28.4	30.5	28.7	24.2	20.4
	Rain	1.2	1.6	0.6	1.0	1.1	0.3	0.2
	SRAD	20.4	23.0	25.9	25.7	21.4	17.3	12.5
Wooster, OH 1988	Tmin	3.8	7.6	10.7	15.9	16.3	10.6	–
	Tmax	16.6	23.1	28.3	32.2	29.0	23.4	–
	Rain	1.9	1.1	0.4	5.2	2.8	2.5	–
	SRAD	15.0	20.3	23.9	20.3	18.0	13.6	–
Wooster, OH 1990	Tmin	7.3	13.2	15.5	14.6	11.2	5.6	–
	Tmax	19.6	25.7	27.4	26.8	22.9	17.7	–
	Rain	4.2	2.0	5.5	3.7	3.7	3.9	–
	SRAD	17.4	20.7	20.1	16.7	13.1	9.5	–
Gainesville, FL 1979	Tmin	15.1	17.0	20.1	22.6	–	–	–
	Tmax	28.2	30.0	32.6	34.0	–	–	–
	Rain	6.9	2.8	3.9	3.6	–	–	–
	SRAD	19.1	21.3	20.4	21.1	–	–	–

† Tmin, minimum temperature (°C); Tmax, maximum temperature (°C); Rain, average daily rainfall (mm); and SRAD, solar radiation (MJ/m²/d).

Results and Discussion

Maize Results

Comparison of Simulated and Observed Crop Variables

A comparison of results between field observed and model simulated output variables of Prisma maize from three different sources of CSPs is summarized in Table 13–5. The GLUE-estimated CSPs did a better job than the arbitrary generic maize cultivar default CSPs and performed as well as the hand-calibrated CSPs in simulating this maize cultivar. The average RAE values were only 3 and 8% in 1995 and 1996, respectively, while they were 4 and 10% for the hand-calibrated CSPs and above 15% for the default CSPs. Using both the GLUE-estimated and hand-calibrated CSPs, the maize model predicted phenology dates very well. In 1995, there were 0 and 2 d differences between the GLUE simulated and observed anthesis and maturity dates respectively, while these values were 1 and 2 d in 1996. Results for the hand-calibrated CSPs were similar. However, the errors in phenology dates simulations with the default CSPs were high. For example, the

Table 13–5. Comparison between observations and model output variables of ‘Prisma’ maize grown at Zaragoza, Spain

Source of outputs	Phenology				Aboveground						
	Anthesis		Maturity		Grain yield		Biomass		Max LAI		Avg.
	Value	RAE†	Value	RAE	Value	RAE	Value	RAE	Value	RAE	RAE
	—————d—————				—————kg/ha—————						
Zaragoza, Spain in 1995											
Field observations	78	–	141	–	10,960	–	23,970	–	6.15	–	–
Default CSPs	62	21%	113	20%	12,679	16%	26,428	10%	5.19	16%	17%
Hand-calibrated CSPs	78	0%	141	0%	10,255	6%	25,648	7%	5.61	9%	4%
GLUE-calibrated CSPs	78	0%	139	1%	10,866	1%	26,437	10%	5.65	8%	3%
Zaragoza, Spain in 1996											
Field observations	78	–	147	–	12,340	–	22,730	–	4.75	–	–
Default CSPs	67	14%	129	12%	13,626	10%	28,973	27%	5.69	20%	16%
Hand-calibrated CSPs	79	1%	151	3%	10,202	17%	25,327	11%	5.55	17%	10%
GLUE-calibrated CSPs	79	1%	149	1%	10,791	13%	26,079	15%	5.56	17%	8%

† The RAE values are the relative absolute error between field observed and model simulated variables from three different sources of CSPs.

anthesis and maturity dates simulated with default CSPs were 11 and 18 d sooner than the field observations in 1996. Although the hand-calibrated CSPs for this cultivar performed about as well as the GLUE-estimated CSPs, it is dangerous for users to automatically use CSPs for a particular cultivar without evaluating them for their own conditions, even if the cultivar’s parameter set is contained in the DSSAT database. Data from regions or locations where the model is to be applied should be used to evaluate simulated results relative to observations or used to estimate a new set of CSPs.

The average absolute error (RAE) between simulated and observed data was lower in 1995 for the GLUE-estimated CSPs (3%) than for the default or hand-calibrated CSPs (17 and 4%, respectively) (Table 13–5). In addition, the RAE values were lower for all but one variable (biomass) when GLUE-estimated parameters were used in 1995 and 1996. In 1996, the GLUE-estimated parameters performed better on average as well; the RAE was 8% in comparison with 16 and 10% for default and hand-calibrated CSPs, respectively.

Comparison of CSPs from GLUE with Hand-Calibrated and Default CSPs

The parameter values of Prisma maize obtained from different sources were compared (Table 13–6). There were large differences among the values of P1 and P2, which determine the thermal time of the vegetative and reproductive stages of maize. Thus, it is not surprising to see large differences among the

Table 13–6. Comparison between cultivar-specific parameters (CSPs) of ‘Prisma’ maize.

Source of CSPs	P1	P2	P5	G2	G3
Default	200.0	0.300	800.0	700.0	8.50
Hand-calibrated	280.0	0.300	789.0	650.0	6.03
GLUE-calibrated values (standard deviations)	268.2 (18.45)	0.758 (0.291)	770.8 (35.72)	675.1 (59.47)	6.439 (0.234)

predicted anthesis and maturity dates with these different sources of CSPs. The standard deviations were mostly less than 10% of the estimated CSPs (Table 13–6). An exception was for the parameter P2, the photoperiod sensitivity coefficient, which was nearly 40% of the estimated value. This is one indication that the model was not sensitive to this CSP for these two experiments, which was apparently due to the fact that the experiments were in the same location both years, and there was little difference in daylengths that the two crops experienced.

Comparing In-Season Simulated Results

Although time series data were not used to estimate CSPs in the GLUE procedure, we show time-series model outputs of in-season predictions for the three sources of CSPs for one season of 1995 (Fig. 13–2). In the maize experiments, no in-season data were collected, so it is not possible to draw conclusions about how well each set of CSPs performed in this example. However, it is interesting to show how different the in-season results are for two reasons. First, it is clear that similar or the same end of season simulations can be obtained with different CSPs, but that considerable differences may occur among in-season results. This can be seen in the GLUE simulated end point results for biomass and grain yield in Fig. 13–2. Second, in-season measurements may be very useful to refine the CSPs so that they simulate the time courses of growth and yield during a season. The differences among the simulated values of leaf area index (LAI) and grain weight (kg ha^{-1}) can be seen clearly.

Soybean Results

Comparison of Simulated and Observed Crop Variables

Similarly, comparisons of the results for Williams soybean are tabulated in Table 13–7 for observed and simulated output variables for the three experiments in two locations. The GLUE and hand-calibrated RAE results averaged across all measurements were nearly equal. In about one-half of the RAE values for individual measurement variables, the GLUE method performed better, and in the other half, the hand-calibrated CSPs performed better. For soybean, Dr. K.J. Boote




Fig. 13–2. Graphs of simulated results using the three sources of cultivar-specific parameters (CSPs) for (a) leaf area index and (b) total biomass and grain yield for the experiment in Zaragoza, Spain in 1995. “Default” represents the outputs from default genetic coefficients, “GLUE-estimated” represents the outputs from GLUE-estimated genetic coefficients, and “Hand-calibrated” represents the outputs from hand-calibrated genetic coefficients.

of the Agronomy Department of the University of Florida (personal communication, 2010) estimated the CSPs using both end of season and phenology data with considerable emphasis on in-season measurements. Thus, the GLUE method did a good job in estimation of CSPs, since the average RAE values were equal to the hand-calibrated CSPs and clearly less than most of the default CSPs. One exception was the maximum LAI (MaxLAI), for which the RAE obtained from the GLUE-estimated CSPs was 31%. The GLUE method also improved the accuracy

Table 13-7. Comparison between observations and model output variables of ‘Williams’ soybean. Default CSPs are those for ‘generic’ MG 4 cultivar.

Source of outputs	Phenology				Aboveground																				
	Anthesis		Maturity		Grain yield			Biomass			Max LAI			First pod			Pod wt.			Unit wt. grain			Avg.		
	Value	RAE†	Value	RAE	Value	RAE	Value	RAE	Value	RAE	Value	RAE	Value	RAE	Value	RAE	Value	RAE	Value	RAE	Value	RAE	Value	RAE	
Wooster, OH, 1988																									
Observed	71	-	145	-	3976	-	8090	-	8090	-	7.65	-	91	-	5194	0%	0.151	-	-	-	-	-	-	-	-
Default CSPs	87	23%	-	-	3138	21%	8356	3%	8356	3%	7.67	0%	101	11%	4490	14%	0.16	6%	11%	11%	6%	11%	6%	11%	11%
Hand-calibrated CSPs	73	3%	141	3%	3781	5%	7892	2%	7892	2%	6.89	10%	90	1%	5145	1%	0.157	4%	4%	4%	4%	4%	4%	4%	4%
GLUE-calibrated CSPs	72	1%	145	0%	3893	2%	7664	5%	7664	5%	6.74	12%	89	2%	5163	1%	0.161	7%	7%	7%	7%	7%	7%	7%	7%
Wooster, OH, 1990																									
Observed	-	-	150	-	3149	-	7113	-	7113	-	7.21	-	96	-	4915	0%	0.164	-	-	-	-	-	-	-	-
Default CSPs	88	-	162	8%	3718	18%	8387	18%	8387	18%	7.47	4%	104	8%	5127	4%	0.168	2%	9%	9%	2%	9%	2%	9%	9%
Hand-calibrated CSPs	73	-	143	5%	3552	13%	7450	5%	7450	5%	6.82	5%	91	5%	4883	1%	0.146	11%	6%	6%	11%	6%	11%	6%	6%
GLUE-calibrated CSPs	71	-	148	1%	3713	18%	7274	2%	7274	2%	6.61	8%	90	6%	4957	1%	0.156	5%	6%	6%	5%	6%	5%	6%	6%
Gainesville, FL, 1979																									
Observed	38	-	93	-	2474	-	3893	-	3893	-	3.15	-	48	-	-	-	0.175	-	0%	0%	-	-	-	-	0%
Default CSPs	40	5%	114	23%	3570	44%	5700	46%	5700	46%	3.38	7%	50	4%	4936	-	0.183	5%	19%	19%	5%	19%	5%	19%	19%
Hand-calibrated CSPs	37	3%	96	3%	2831	14%	4303	11%	4303	11%	2.55	19%	48	0%	3893	-	0.156	11%	9%	9%	11%	9%	11%	9%	9%
GLUE-calibrated CSPs	37	3%	99	6%	2588	5%	3855	1%	3855	1%	2.16	31%	47	2%	3416	-	0.169	3%	7%	7%	3%	7%	3%	7%	7%

† The RAE values are the relative absolute error between field observed and model simulated variables from three different sources of CSPs.

of phenology date predictions relative to the other sources of CSPs. The differences between observed and GLUE simulated anthesis and maturity dates were all less than 2 d, except for the maturity date of the experiment in Gainesville, FL, in 1979, which was about 6 d.

Comparison of CSPs from GLUE with Hand-Calibrated and Default CSPs

The parameter values of Williams soybean obtained from different sources are compared in Table 13–8. The default CSPs were assumed to be those of a generic maturity 4 soybean variety from the DSSAT database. Except for parameters SLAVR and SIZELF, the differences between these three types of CSPs were very small. The largest relative differences between hand-calibrated and GLUE-estimated CSPs were for PPSEN (0.285 vs. 0.342, respectively). The values of CSDL and PPSEN for Williams in hand-calibrated CSP were not hand-calibrated, in fact, but were estimated using a least squares simplex method from a much larger ($n > 100$) data set (Grimm et al., 1993). Thus, the GLUE-estimated photoperiod sensitivity coefficient (PPSEN) was obtained from a much smaller set of environments and is less robust across regions than that obtained by Grimm et al. (1993) for the hand-calibrated CSPs. This highlights a fundamental issue that transcends methods for calibration—that CSPs estimated from field data may not be robust if the size of the dataset used is small. Other relatively large differences occurred for SLAVR and SIZELF. Nevertheless, the standard deviations of the GLUE-estimated CSPs were relatively low, indicating that the parameters worked well across the locations and years for this soybean cultivar.

Comparison of In-Season Simulated and Observed Soybean Results

The time-series model outputs were also compared for Williams soybean cultivar (Fig. 13–3, 13–4, 13–5). All three experiments in this study provided field observations for some time-series output variables, such as LAI, biomass, and grain weight. These observations can help us to evaluate the reliability of the predictions from different CSPs sources. Most model outputs simulated using the GLUE-estimated and hand-calibrated CSPs were similar for all variables that were measured during the growing season. This result provides strong evidence that the GLUE method provided reliable CSP estimates, similar in performance to those that were estimated by an expert. However, this result also demonstrates the concept of “equifinality” (Beven and Freer, 2001; Hansson and Lundin, 2006; Shulz et al., 1999), which means that different combinations of CSPs will result in the same quality of simulation. The values of the GLUE CSPs were different from those estimated by K.J. Boote, yet simu-

Table 13–8. Comparison among cultivar-specific parameters (CSPs) of ‘Williams’ soybean. Default CSPs are those for “generic” MG 4 cultivar.

Source of CSPs	CSDL	PPSEN	EM-FL	FL-SD	SD-PM	LFMAX	SLAVR	SIZELF	WTSPD	SFDUR	SDPDV
Default	13.09	0.294	19.4	15.0	34.00	1.030	375.0	180.0	0.190	23.0	2.20
Hand-calibrated	13.40	0.285	19.0	13.8	32.20	1.000	385.0	180.0	0.180	26.0	2.40
GLUE-calibrated	13.67	0.342	18.7	15.4	36.23	1.004	425.0	143.1	0.174	23.4	2.23
values (SD)	(0.384)	(0.059)	(1.206)	(1.132)	(2.209)	(0.025)	(15.60)	(13.98)	(0.008)	(1.445)	(0.21)




Figure 13–3. Graphs of observed and simulated results using the three sources of cultivar-specific parameters (CSPs) for (a) leaf area index and (b) total biomass and grain yield for the experiment in Wooster, OH in 1988. “Default” represents the outputs from default genetic coefficients, “GLUE-estimated” represents the outputs from GLUE-estimated genetic coefficients, and “Hand-calibrated” represents the outputs from hand-calibrated genetic coefficients by K.J. Boote.

lated results and errors in prediction were nearly equal showing that different “best” CSP sets might exist. However, the GLUE-estimated CSPs did not correctly predict LAI for the Gainesville, FL 1979 experiment. It is not surprising to see that the GLUE method did not improve time-series prediction because there were no time-series observations involved in estimating the CSPs in the GLUE approach used in this study.




Figure 13–4. Graphs of observed and simulated results using the three sources of cultivar-specific parameters (CSPs) for (a) leaf area index and (b) total biomass and grain yield for the experiment in Wooster, OH in 1990. “Default” represents the outputs from default genetic coefficients, “GLUE-estimated” represent the outputs from GLUE-estimated genetic coefficients, and “Hand-calibrated” represent the outputs from hand-calibrated genetic coefficients by K.J. Boote.

Conclusions

The use of field observations to estimate CSPs of cropping system models is necessary for practical uses in predicting crop performance under different soil, climate, and management scenarios. Various methods can be used to estimate CSPs, each with advantages and disadvantages. We presented the GLUE method in this chapter that has been implemented for estimating CSPs for all crops in




Figure 13-5. Graphs of observed and simulated results using the three sources of cultivar-specific parameters (CSPs) for (a) leaf area index and (b) total biomass and grain yield for the experiment in Gainesville, FL in 1979. “Default” represents the outputs from default genetic coefficients, “GLUE-estimated” represent the outputs from GLUE-estimated genetic coefficients, and “Hand-calibrated” represent the outputs from hand-calibrated genetic coefficients by K.J. Boote.

DSSAT v4.5, using field observations made once during a season. We showed that this method works well for two of the crops in DSSAT. However, there are important theoretical considerations that users should be aware of when using any method to estimate CSPs using field data. First, any calibration process may result in CSPs that are not generally applicable, particularly if the range of environments in which data were collected is narrow or if the crops experience drought, nutrient, or other stresses that are not adequately simulated by the model. In the

first situation, having a narrow range of environments may result in parameters that work well only for those environments. For example, if datasets with a narrow range of daylengths are used, the CSPs that determine photoperiod response in soybean (CSDL and PPSN) are not likely to be robust for use across locations with daylengths outside that range. In the second situation, it is likely that the estimation process will attempt to set CSP values to compensate for water, nutrient, or pest stresses if they occur and are not adequately simulated in the model. This problem can happen when the model does not include mechanisms for simulating those stresses, or when the soil parameters or initial conditions are not accurate. This is a main reason for recommending the use of irrigated and well-fertilized treatments for estimating CSPs. This is true for the GLUE and GENCALC methods available in DSSAT as well as any other procedure where the two described situations occur. Another principle that is important regardless of method used is that the model itself may not adequately represent the crop development and growth processes adequately. In this case, one may obtain estimates from a particular set of field data that work well for that set of data but may not be robust. Thus, it is important for users to evaluate any existing CSPs for their own conditions if they were estimated in other environments.

In this chapter, two cultivars were selected, one for each of two crops (maize and soybean), to evaluate the performance of GLUE program. It was shown that the GLUE-estimated CSPs for maize were different from the default values and from the manually estimated parameters. Similarly, the soybean CSPs estimated by the GLUE procedure were different from the arbitrary default set. However, soybean CSPs were similar to those manually estimated by an expert model user. Simulated outputs using the GLUE-estimated CSPs and hand-calibrated CSPs were similar to observed maize values. Both compared more favorably to observed maize values than those obtained from the default CSPs. For soybean, the simulated outputs were superior to the default set and equivalent to those obtained using the hand-calibrated CSPs. The average RAE values were all smaller than those for the default set and generally equal to those for the hand-calibrated CSPs in this study. There are limitations of the GLUE program as implemented in DSSAT. First, it does not include time-series measurements. As shown in the comparisons with time-series observations, the GLUE method did not always improve the accuracy of in-season model predictions, but it also did not deteriorate accuracy. This limitation can be overcome in future implementations of methods to estimate CSPs. Second, the GLUE program requires a large number of model runs, which may require more than 1 h on modern microcomputers. Nevertheless, implementation of the GLUE procedure in DSSAT v4.5 provides users an easy to use option for estimating CSPs from field observations for cultivars grown in their region.

Acknowledgments

This contribution was partially supported by the Office of Natural Resources Management and Office of Agriculture in the Economic Growth, Agriculture, and Trade Bureau of the U.S. Agency for International Development, under terms of Grant No. LAG-G-00-97-00002-00. Special appreciation is expressed to Dr. Michael D. Dukes of the Department of Agricultural and Biological Engineering, University of Florida for his help on this project.

References

- Aronica, G., P.D. Bates, and M.S. Horritt. 2002. Assessing the uncertainty in distributed model predictions using observed binary pattern information within GLUE. *Hydrol. Processes* 16:2001–2016.
- Baldi, P.F., M.C. Vanier, and J.M. Bower. 1998. On the use of Bayesian methods for evaluating compartmental neural models. *J. Comput. Neurosci.* 5:285–314.
- Bannayan, M., and G. Hoogenboom. 2008. Weather analog: A tool for real-time prediction of daily weather data realizations based on a modified K-nearest neighbor approach. *Environ. Model. Softw.* 23:703–713.
- Bertin, N., P. Martre, M. Génard, B. Quilot, and C. Salon. 2009. Under what circumstances can process-based simulation models link genotype to phenotype for complex traits? Case study of fruit and grain quality traits. *J. Exp. Bot.* 61:955–967, doi:10.1093/jxb/erp377.
- Beven, K., and A. Binley. 1992. The future of distributed models: Model calibration and uncertainty prediction. *Hydrol. Processes* 6:279–298.
- Beven, K., and J. Freer. 2001. Equifinality, data assimilation, and uncertainty estimation in mechanistic modeling of complex environmental systems using the GLUE methodology. *J. Hydrol.* 249:11–29.
- Bhalla, U.S., and J.M. Bower. 1993. Exploring parameter space in detailed single neuron models: Simulations of the mitral and granule cells of the olfactory-bulb. *J. Neurophysiol.* 69:1948–1965.
- Blasone, R.S., J.A. Vrugt, H. Madsen, D. Rosbjerg, B.A. Robinson, and G.A. Zyvoloski. 2008. Generalized likelihood uncertainty estimation (GLUE) using adaptive Markov Chain Monte Carlo sampling. *Adv. Water Resour.* 31:630–648.
- Boote, K.J. 1999. Concepts for calibrating crop growth models. p. 179–200. *In* G. Hoogenboom, P.W. Wilkens, and G.Y. Tsuji (ed.) DSSAT Version 3. A Decision Support system for Agrotechnology Transfer. Vol. 4. University of Hawaii, Honolulu.
- Boote, K.J., J.W. Jones, W.D. Batchelor, E.D. Nafziger, and O. Myers. 2003. Genetic coefficients in the CROPGRO–Soybean model: Links to field performance and genomics. *Agron. J.* 95:32–51.
- Boote, K.J., M.J. Kropff, and P.S. Bindraban. 2001. Physiology and modeling of traits in crop plants: Implications for genetic improvement. *Agric. Syst.* 70:395–420.
- Brazier, R.E., K.J. Beven, S.G. Anthony, and J.S. Rowan. 2001. Implications of model uncertainty for the mapping of hillslope-scale soil erosion predictions. *Earth Surf. Process. Landf.* 26:1333–1352.
- Campbell, E.P., D.R. Fox, and B.C. Bates. 1999. A Bayesian approach to parameter estimation and pooling in nonlinear flood event models. *Water Resour. Res.* 35:211–220.
- Eichler-West, R., and G. Wilcox. 1997. Robust parameter selection for compartmental models of neurons using evolutionary algorithms. *In* J.M. Bower (ed.) *Computational neuroscience: Trends in research 1997*. Plenum Publishing, New York.
- Feyen, L., K.J. Beven, F. De Smedt, and J. Freer. 2001. Stochastic capture zone delineation within the generalized likelihood uncertainty estimation methodology: Conditioning on head observations. *Water Resour. Res.* 37:625–638.
- Foster, W.R., L.H. Ungar, and J.S. Schwaber. 1993. Significance of conductances in Hodgkin-Huxley models. *J. Neurophysiol.* 70:2502–2518.
- Franks, S.W., K.J. Beven, P.F. Quinn, and I.R. Wright. 1997. On the sensitivity of soil–vegetation–atmosphere transfer (SVAT) schemes: Equifinality and the problem of robust calibration. *Agric. For. Meteorol.* 86:63–75.

- Franks, S.W., P. Gineste, K.J. Beven, and P. Merot. 1998. On constraining the predictions of a distributed model: The incorporation of fuzzy estimates of saturated areas into the calibration process. *Water Resour. Res.* 34:787–797.
- Freer, J., K.J. Beven, and B. Ambroise. 1996. Bayesian estimation of uncertainty in runoff prediction and the value of data: An application of the GLUE approach. *Water Resour. Res.* 32:2161–2173.
- Grimm, S.S., J.W. Jones, K.J. Boote, and J.D. Hesketh. 1993. Parameter estimation for predicting flowering date of soybean cultivars. *Crop Sci.* 33:137–144.
- Hankin, B.G., R. Hardy, H. Kettle, and K.J. Beven. 2001. Using CFD in a GLUE framework to model the flow and dispersion characteristics of a natural fluvial dead zone. *Earth Surf. Process. Landf.* 26:667–687.
- Hansson, K., and C. Lundin. 2006. Equifinality and sensitivity in freezing and thawing simulations of laboratory and in situ data. *Cold Reg. Sci. Technol.* 44:20–37.
- He, J. 2008. Best management practice development with the CERES-Maize model for sweet corn production in North Florida. Ph.D. diss. Agricultural and Biological Engineering Dep., University of Florida, Gainesville.
- He, J., M.D. Dukes, J.W. Jones, W.D. Graham, and J. Judge. 2009. Applying GLUE for estimating CERES-Maize genetic and soil parameters for sweet corn production. *Trans. ASABE* 52:1907–1921.
- He, J., J.W. Jones, W.D. Graham, and M.D. Dukes. 2010. Influence of likelihood function choice for estimating crop model parameters using the generalized likelihood uncertainty estimation method. *Agric. Syst.* 103:256–264, doi:10.1016/j.agsy.2010.01.006.
- Hoogenboom, G., J.W. Jones, C.H. Porter, P.W. Wilkens, K.J. Boote, W.D. Batchelor, L.A. Hunt, and G.Y. Tsuji. 2003. Decision Support System for Agrotechnology Transfer Version 4.0 [CD]. Univ. of Hawaii, Honolulu.
- Hoogenboom, G., J.W. Jones, P.W. Wilkens, C.H. Porter, K.J. Boote, L.A. Hunt, U. Singh, J.L. Lisazo, J.W. White, O. Uryasev, F.S. Royce, R. Ogoshi, A.J. Gijsman, and G.Y. Tsuji. 2010. Decision Support System for Agrotechnology Transfer Version 4.5 [CD]. Univ. of Hawaii, Honolulu.
- Hunt, L.A., S. Pararajasingham, J.W. Jones, G. Hoogenboom, D.T. Imamura, and R.M. Ogoshi. 1993. GenCalc: Software to facilitate the use of crop models for analyzing field experiments. *Agron. J.* 85:1090–1094.
- Jawitz, J.W., R. Muñoz-Carpena, S. Muller, K.A. Grace, and A.I. James. 2008. Development, Testing, and Sensitivity and Uncertainty Analyses of a Transport and Reaction Simulation Engine (TaRSE) for Spatially Distributed Modeling of Phosphorus in South Florida Peat Marsh Wetlands. Scientific Investigations Rep. 2008-5029. USGS, Reston, VA.
- Jensen, J.B. 2003. Parameter and uncertainty estimation in groundwater modeling. Ph.D. diss. Series Paper 23. Dep. of Civil Engineering, Aalborg Univ., Denmark.
- Jones, J.W., G. Hoogenboom, C. Porter, K.J. Boote, W.D. Batchelor, L.A. Hunt, P. Wilkens, U. Singh, A. Gijsman, and J.T. Ritchie. 2003. The DSSAT model cropping system model. *Eur. J. Agron.* 18:235–265.
- Jones, J.W., B.A. Keating, and C.H. Porter. 2001. Approaches to modular model development. *Agric. Syst.* 70:421–443.
- Jones, J.W., J.B. Naab, D. Fatondji, K. Dzotsi, S. Adiku, and J. He. 2010. Uncertainties in simulating crop performance in degraded soils and low input production systems. p. 218–219. *In* M. van Ittersum et al. (ed.) *Integrated assessment of agricultural and sustainable development: Setting the agenda for science and policy. (AgSAP 2009)*, 10–12 Mar. 2009. Egmond aan Zee, The Netherlands.
- Jones, J.W., G.Y. Tsuji, G. Hoogenboom, L.A. Hunt, P.K. Thornton, P.W. Wilkens, D.T. Imamura, W.T. Bowen, and U. Singh. 1998. Decision support system for agrotechnology transfer; DSSAT v3. p. 157–177. *In* G.Y. Tsuji et al. (ed.) *Understanding options for agricultural production*. Kluwer Academic Publ., Dordrecht, the Netherlands.
- Lamb, R., K. Beven, and S. Myrab. 1998. Use of spatially distributed water table observations to constrain uncertainty in a rainfall–runoff model. *Adv. Water Resour.* 22:305–317.
- Lyneis, J.M., and A.L. Pugh. 1996. Automated vs. hand calibration of system dynamics models: An experiment with a simple project model. p. 317–320. *In* G.P. Richardson and J.D. Sterman (ed.) *Proceedings of the 1996 International System Dynamics Conference*. System Dynamics Society, Cambridge, MA.
- Madsen, H., G. Wilson, and H.C. Ammentorp. 2002. Comparison of different automated strategies for calibration of rainfall–runoff models. *J. Hydrol.* 261:48–99.

- Makowski, D., J. Hillier, D. Wallach, B. Andrieu, and M.H. Jeuffroy. 2006. Parameter estimation for crop models. p. 101–150. *In* D. Wallach et al. (ed.) *Working with dynamic crop models*. Elsevier, Amsterdam.
- Makowski, D., D. Wallach, and M. Tremblay. 2002. Using a Bayesian approach to parameter estimation: Comparison of the GLUE and MCMC methods. *Agronomie* 22:191–203.
- Malakoff, D. 1999. Bayes offers a 'New' way to make sense of numbers. *Science* 286:1460–1464.
- Mavromatis, T., K.J. Boote, J.W. Jones, A. Irmak, D. Shinde, and G. Hoogenboom. 2001. Developing genetic coefficients for crop simulation models with data from crop performance trials. *Crop Sci.* 41:40–51.
- Mavromatis, T., K.J. Boote, J.W. Jones, G.G. Wilkerson, and G. Hoogenboom. 2002. Repeatability of model genetic coefficients derived from soybean performance trials across different states. *Crop Sci.* 42:76–89.
- McMichael, C.E., A.S. Hope, and H.A. Loaiciga. 2006. Distributed hydrological modeling in California semiarid shrublands: MIKE SHE model calibration and uncertainty estimation. *J. Hydrol.* 317:307–324.
- Mertens, J., H. Madsen, L. Feyen, D. Jacques, and J. Feyen. 2004. Including prior information in the estimation of effective soil parameters in unsaturated zone modeling. *J. Hydrol.* 294:251–269.
- Monod, H., C. Naud, and D. Makowski. 2006. Uncertainty and sensitivity analysis for crop models. p. 55–99. *In* D. Wallach et al. (ed.) *Working with dynamic crop models*. Elsevier, Amsterdam.
- Muleta, M.K., and J. Nicklow. 2005. Sensitivity and uncertainty analysis coupled with automatic calibration for a distributed watershed model. *J. Hydrol.* 306:127–145.
- Müller, T., J. Magid, L.S. Jensen, and N.E. Nielsen. 2003. Decomposition of plant residues of different quality in soil—DAISY model calibration and simulation based on experimental data. *Ecol. Model.* 166:3–18.
- Muñoz-Carpena, R., G.A. Fox, and G.J. Sabbagh. 2010. Parameter importance and uncertainty in predicting runoff pesticide reduction with filter strips. *J. Environ. Qual.* 39:1–12, doi:10.1016/10.2134/jeq2009.0300.
- Muñoz-Carpena, R., Z. Zajac, and Y.-M. Kuo. 2007. Evaluation of water quality models through global sensitivity and uncertainty analyses techniques: Application to the vegetative filter strip model VFSSMOD-W. *Trans. ASABE* 50:1719–1732.
- Oliva, R. 2003. Model calibration as a testing strategy for system dynamics models. *Eur. J. Oper. Res.* 151:552–568.
- Porter, C., J.W. Jones, and R. Braga. 2000. An approach for modular crop model development. International Consortium for Agricultural Systems Applications, Honolulu, HI.
- R Development Core Team. 2009. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Ritchie, J.T., J.R. Kiniry, C.J. Jones, and P.T. Dyke. 1986. Model inputs. p. 37–48. *In* C.A. Jones and J.R. Kiniry (ed.) *CERES-Maize: A simulation model of maize growth and development*. Texas A&M Univ. Press, College Station, TX.
- Romanowicz, R., and K.J. Beven. 2006. Comments on generalized likelihood uncertainty estimation. *Reliab. Eng. Syst. Saf.* 91:1315–1321.
- Romanowicz, R., K.J. Beven, and J. Tawn. 1994. Evaluation of predictive uncertainty in nonlinear hydrological models using a Bayesian approach. p. 297–317. *In* V. Barnett and K.F. Turkman (ed.) *Statistics for the environment II. Water related issues*. John Wiley and Sons, New York.
- Romanowicz, R.J., K.J. Beven, and J. Tawn. 1996. Bayesian calibration of flood inundation models. p. 333–360. *In* M.G. Anderson et al. (ed.) *Floodplain processes*. John Wiley and Sons, New York.
- Saltelli, A., S. Tarantola, F. Campolongo, and M. Ratto. 2004. *Sensitivity analysis in practice: A guide to assessing scientific models*. John Wiley and Sons, Chichester, UK.
- Shulz, K., K. Beven, and B. Huwe. 1999. Equifinality and the problem of robust calibration in nitrogen budget simulations. *Soil Sci. Soc. Am. J.* 63:1934–1941.
- Tadesse, A., and E.N. Anagnostou. 2005. A statistical approach to ground radar–rainfall estimation. *J. Atm. Ocean Technol.* 22:1055–1071.
- Timsina, J., and E. Humphreys. 2006. Performance of CERES-Rice and CERES-Wheat models in rice–wheat system: A review. *Agric. Syst.* 90:5–31.
- Tsujii, G.Y., G. Uehara, and S. Balas. 1994. DSSAT V.3. University of Hawaii, Honolulu.

- Uehara, G. 1998. Synthesis. p. 389–392. *In* G.Y. Tsuji et al. (ed.) Understanding options for agricultural production. Kluwer Academic Publ., Dordrecht, The Netherlands.
- Vanier, M., and J. Bower. 1996. A comparison of automated parametersearching methods for neural models. *In* J.M. Bower (ed.) Proceedings of the 1995 Computational Neuroscience Conference (CNS*95), Monterey (CA). Academic Press, New York.
- Wallach, D., B. Goffinet, J.E. Bergez, P. Debaeke, D. Leenhardt, and J.N. Aubertot. 2001. Parameter estimation for crop models: A new approach and application to a corn model. *Agron. J.* 93:757–766.
- Wang, X., X. He, J.R. Williams, R.C. Izaurralde, and J.D. Atwood. 2005. Sensitivity and uncertainty analyses of crop yields and soil organic carbon simulated with EPIC. *Trans. ASAE* 48:1041–1054.