

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/268211838>

Development of soil and terrain digital database for major food-growing regions of India for resource planning

Article in *Current Science* · November 2014

CITATIONS

21

READS

1,018

87 authors, including:



P. Chandran

198 PUBLICATIONS 3,785 CITATIONS

SEE PROFILE



Pramod Tiwary

National Bureau of Soil Survey and Land Use Planning

136 PUBLICATIONS 1,723 CITATIONS

SEE PROFILE



Tapas Bhattacharyya

Dr. Balasaheb Sawant Konkan Krishi Vidypeeth

303 PUBLICATIONS 6,223 CITATIONS

SEE PROFILE



Champa Mandal

National Bureau of Soil Survey and Land Use Planning

75 PUBLICATIONS 2,127 CITATIONS

SEE PROFILE

Development of soil and terrain digital database for major food-growing regions of India for resource planning

P. Chandran^{1*}, P. Tiwary¹, T. Bhattacharyya¹, C. Mandal¹, J. Prasad¹, S. K. Ray¹, D. Sarkar¹, D. K. Pal², D. K. Mandal¹, G. S. Sidhu³, K. M. Nair⁴, A. K. Sahoo⁵, T. H. Das⁵, R. S. Singh⁶, R. Srivastava¹, T. K. Sen¹, S. Chatterji¹, N. G. Patil¹, G. P. Obireddy¹, S. K. Mahapatra³, K. S. Anil Kumar⁴, K. Das⁵, A. K. Singh⁶, S. K. Reza⁷, D. Dutta⁵, S. Srinivas⁴, K. Karthikeyan¹, M. V. Venugopalan⁸, K. Velmourougane⁸, A. Srivastava⁹, Mausumi Raychaudhuri¹⁰, D. K. Kundu¹⁰, K. G. Mandal¹⁰, G. Kar¹⁰, J. A. Dijkshoorn¹¹, N. H. Batjes¹¹, P. S. Bindraban¹¹, S. L. Durge¹, G. K. Kamble¹, M. S. Gaikwad¹, A. M. Nimkar¹, S. V. Bobade¹, S. G. Anantwar¹, S. V. Patil¹, K. M. Gaikwad¹, V. T. Sahu¹, H. Bhondwe¹, S. S. Dohtre¹, S. Gharami¹, S. G. Khapekar¹, A. Koyal⁴, K. Sujatha⁴, B. M. N. Reddy⁴, P. Sreekumar⁴, D. P. Dutta⁷, L. Gogoi⁷, V. N. Parhad¹, A. S. Halder⁵, R. Basu⁵, R. Singh⁶, B. L. Jat⁶, D. L. Oad⁶, N. R. Ola⁶, K. Wadhai¹, M. Lokhande¹, V. T. Dongare¹, A. Hukare¹, N. Bansod¹, A. H. Kolhe¹, J. Khuspure¹, H. Kuchankar¹, D. Balbuddhe¹, S. Sheikh¹, B. P. Sunitha⁴, B. Mohanty³, D. Hazarika⁷, S. Majumdar⁵, R. S. Garhwal⁶, A. Sahu⁸, S. Mahapatra¹⁰, S. Puspamitra¹⁰, A. Kumar⁹, N. Gautam¹, B. A. Telpande¹, A. M. Nimje¹, C. Likhar¹ and S. Thakre¹

¹Regional Centre, National Bureau of Soil Survey and Land Use Planning, Nagpur 440 033, India

²International Crops Research Institute for the Semi-Arid Tropics, Patancheru 502 324, India

³Regional Centre, National Bureau of Soil Survey and Land Use Planning, New Delhi, 110 012, India

⁴Regional Centre, National Bureau of Soil Survey and Land Use Planning, Bangalore 560 024, India

⁵Regional Centre, National Bureau of Soil Survey and Land Use Planning, Kolkata 700 091, India

⁶Regional Centre, National Bureau of Soil Survey and Land Use Planning, Udaipur 313 001, India

⁷Regional Centre, National Bureau of Soil Survey and Land Use Planning, Jorhat 785 004, India

⁸Central Institute for Cotton Research, Nagpur 440 010, India

⁹National Bureau of Agriculturally Important Microorganisms, Mau 275 101, India

¹⁰Directorate of Water Management, Bhubaneswar 751 023, India

¹¹ISRIC, Wageningen, The Netherlands

Soil information system in SOTER (soil and terrain digital database) framework is developed for the Indo-Gangetic Plains (IGP) and black soil regions (BSR) of India with the help of information from 842 georeferenced soil profiles including morphological, physical and chemical properties of soils in addition to the site characteristics and climatic information. The database has information from 82 climatic stations that can be linked with the other datasets. The information from this organized database can be easily retrieved for use

and is compatible with the global database. The database can be updated with recent and relevant data as and when they are available. The database has many applications such as inputs for refinement of agro-ecological regions and sub-regions, studies on carbon sequestration, land evaluation and land (crop) planning, soil erosion, soil quality, carbon and crop modelling and other climate change related research. This warehouse of information in a structured framework can be used as a data bank for posterity.

Keywords: Black soil region, database, Indo-Gangetic Plains, SOTER.

Introduction

INNOVATIVE methods are increasingly important to utilize existing soil information and in this context spatial soil information systems play an important role^{1,2}. Soil is an important component of land use planning as it acts both

as a source and sink of energy for many functions of the land. In general, all living and non-living things on earth get their energy for functioning from the soil in the form of nutrients, water and air. In the last 2–3 decades, soil information has become increasingly important to many disciplines to address the conflicting pressure on limited land resources. In addition to farming community, civil engineers and agricultural engineers, environmentalists, urban planners, disaster managers and policy makers also need soil information^{3–5}.

This recent spurt in demand for development of information systems on natural resources is primarily due to the widespread awareness of the need for protection and

*For correspondence. (e-mail: pchandran1960@yahoo.co.in)

preservation of the natural environment. Thus, worldwide there is a renewed awareness on the need for information about soils in digital format to support efficient and wise use of this critical resource. Many countries are now focusing on updating and modernizing their soil resource databases⁶. In this renewed effort the role of geoinformatics has become central not only to storing the data, but also for a range of analytical tasks like manipulating and transforming basic data into a variety of quantitative soil information according to user requirement⁷.

A wide range of soil information in India is available in scattered and unorganized format, but the modern-day information system of any natural resource requires its location in terms of time and space and exact referencing or georeferencing of important spots. Therefore, soil information with exact coordinates can be used for developing a such system. A geographic information system (GIS) is an important tool for georeferencing soil information system (GeoSIS).

The Indo-Gangetic Plains (IGP) as a food bowl of India, produces nearly 50% of the total food grains of the country. During the recent past, including the green revolution and beyond, the IGP has been subjected to major agricultural intensification and high population pressure; consequently, there are reports of decline in productivity and fertility due to adverse changes in some dynamic soil properties and overexploitation of available resources⁸. Contrary to this, the black soil region (BSR) is underutilized, primarily due to its inherent nature of the soil, climate and management-related constraints. The major part of this area is rainfed and climate varies from arid to sub-humid (dry). Therefore, agriculture in this region depends on rainwater storage and release characteristics of soils. There are reports of development of subsoil sodicity in some parts of the arid and semi-arid regions due to the development of pedogenic carbonates⁹, while in some other areas there are reports of occurrence of the palygorskite, a mineral which on irrigation develops a net-like structure and retards water movement¹⁰. In contrast, natural modifiers such as zeolites and gypsum are a boon to farmers as they protect the soils against degradation by modifying hydraulic properties even in the presence of sodicity and palygorskite¹¹⁻¹³.

As mentioned earlier, although the information on soils is huge, it is scattered and therefore needs to be archived in a standardized format to provide georeferenced information in a spatial and digital domain for researchers, policy makers and other managers and users of natural resources. With this in mind, the present article describes the process of developing a georeferenced soil information system in SOTER (soil and terrain digital database) framework and is restricted to the IGP and BSR. The database developed can be utilized for resource planning, in general and for agriculture planning and activities, in particular.

Materials and methods

Methods

The global and national soil and terrain digital databases (SOTER) input software was developed at the International Soil Reference and Information Centre (ISRIC), The Netherlands in collaboration with other international organization, viz. FAO, UNEP and IUSS to create and maintain a global digitized map unit and attributes. The SOTER concept is based on the relationship between the physiography, parent material and soils within an area. It provides data for improved mapping and monitoring of changes in soil and terrain resources. The methodology allows mapping and characterization of areas of land with a distinctive, often repetitive, pattern of landform, lithology, surface form, slope, parent material and soils¹⁴. The collated information is stored in the SOTER framework which is linked to a GIS, permitting a wide range of applications^{8,15-18}. The database allows periodic updating by removing obsolete or irrelevant data. SOTER is a combination of geographical and attribute data. Terrain information is a geographic component indicating the topology of SOTER unit and attribute data give spatial unit characteristics stored in a set of relational database management system (RDBMS) files. A SOTER unit is made by combining information on terrain and soil attributes (Figure 1). The database can also store climatic data, data sources, land use and other auxiliary data which are useful for many other land users. The basic data required for the construction of a SOTER unit are topographic, geomorphological, geological and soil map ideally at the scale of 1 : 250,000 to 1 : 1 m as layers, accompanied by sufficient analytical data for soil characterization and mapping.

In SOTER, the units are given unique identification codes. In the attribute tables for terrain, terrain component and soil component, this identification code is completed with sub-codes for the terrain component and soil component. Both the above attributes are derived from




Figure 1. SOTER attribute database structure with area and point data (1: M = one to many, M: 1 = many to one relations)¹⁴.

Table 1. Map details of the Indo-Gangetic Plains and black soil region*

Region	Source map-ID	Map title	Year	Scale (million)	Minimum latitude	Minimum longitude	Maximum latitude	Maximum longitude	UTM zone	Geodetic datum	Type of source map
India	Nil	Soils of India	2002	1 : 1	6°45'	67.7°	37°6'	97°25'	42–47 N	WGS 1984	Soil map
IGP	IN001	Soils of IGP	2010	1 : 1	21°30'	73°52'	32°15'	92°6'	43–45 N	WGS 1984	Soil map
BSR	IN002	Soils of BSR	2010	1 : 1	8°28'	68°24'	27°1'	89°2'	42–45 N	WGS 1984	Soil map

*Source: Refs 19 and 20. UTM, Universal Trans Marketeter.

the site characteristics of each map unit. The soil component information is stored in three tables, viz. soil component, profile and horizon table. The profile and horizon tables hold attribute data for each profile with the exact location, morphological and laboratory data of each horizon and details of the laboratory. The SOTER structure has a link with each table in the database using primary keys. The database also stores information on laboratory methods followed for analysis and sources of information used for compilation of SOTER.

Materials

Geographic database for the IGP: Earlier, the National Bureau of Soil Survey and Land Use Planning (NBSS&LUP) developed a soil map of India on a 1 : 1 m scale¹⁹ with distinct map units with information on landform, lithology, surface form, parent material and soil. Details of the source map developed for SOTER-IGP and BSR are given in Table 1. The soil map of IGP was derived from the 1 : 1 m soil map of India and this was used as a geographic database (Figure 2) for developing the SOTER IGP²⁰. The IGP is situated between Gurdaspur district (Punjab) in the west and Jalpaiguri district (West Bengal) in the east, and to West Tripura district (Tripura) in the northeast extending from 21°45' to 31°30'N lat. and 74°15' to 91°30'E long. A recently revised area estimate of the IGP is 52.01 m ha (ref. 20). The plain is subdivided into 8 agro-ecoregions (AERs), 17 agro-ecological sub-regions (AESRs) and 6 bioclimatic regions depending upon major physiography, climate and length of the growing period^{21–23}. The IGP has a nearly level physiography (plains) and the parent material is derived from the alluvial deposits of the river Ganga and its tributaries. Therefore, each soil map unit is considered as a separate terrain.

SOTER IGP has 348 map units showing association of soils as dominant and subdominant. As SOTER is a global database, the dominant soils occurring in the terrain are considered as its major soils. Soil attribute data were developed from profiles selected from the master database which contains information on 437 soil profiles^{24,25}, of which only 417 points were georeferenced in the map due to scale limitations. This point information was collated from different sources (Table 2) from the published reports and literature.




Figure 2. Georeferenced soil map of the Indo-Gangetic Plains developed for SOTER IGP.

Geographic database for BSR: The geometric database for SOTER BSR was derived from the 1 : 1 m soil map of India¹⁹. The revision of the BSR boundary was made using ASTER data and georeferenced soil information. The revised total area is 76.4 m ha (Figure 3), which is spread across in 36 AESRs of the country²¹. The BSR has 290 map units distributed mainly in eight states with some sporadic occurrence in non-traditional areas²⁶. Each map unit in the BSR is considered as a separate terrain, as these soil map units were made taking into consideration the major physiography, landform and geology in addition to soil. The attribute database for SOTER BSR was developed by selecting representative profiles from the 448 profiles developed for this region (Table 3), of which 425 points were georeferenced in the map.

Procedure for development of SOTER IGP and SOTER BSR

Development of SOTER database: The attribute data table for terrain, terrain component and soil component

Georeferenced SIS for agricultural LUP

Table 2. Distribution of point data in states, agro-ecological regions (AERs), agro-ecological sub-regions (AESRs) and bioclimatic regions of IGP

State	No. of soil series	AERs	AESRs	Bioclimate*	Source
Punjab	53	2, 4, 9,14	2.1, 2.3, 4.1, 9.1, 14.2	SAd, SAm, A	44–47
Haryana	10	2, 4, 9,14	2.3, 4.1, 9.1, 14.2	SAd, SAm, A	45–47
Uttarakhand	1	9	14.5	SAm	47
Delhi	3	4	4.1	SAd	45, 46
Uttar Pradesh	177	4, 9, 13	4.1, 4.3, 4.4, 9.1, 9.2, 13.1, 13.2	SAm, SHd, SAd	48–57
Bihar	86	9, 13	9.2, 13.1	SHm, SHd	45–47, 57, 58
West Bengal	104	12, 13, 15, 16, 18	12.3, 13.1, 15.1, 15.3, 16.1, 16.2, 18.5	PH, SHm	24, 45–47, 49, 59, 60
Tripura	3	15, 17,	15.3, 17.2	PH	23, 24
Total	437 (417)**	10	17		

*PH, Perhumid; SHm, Sub-humid moist; SHd, Sub-humid dry; Sam, Semi-arid moist; SAd, Semi-arid dry; A, arid.

**No. of points georeferenced (many points overlapped due to scale limitation).




Figure 3. Georeferenced soil map of black soil region developed for SOTER BSR.

for the IGP and BSR were developed from the map and series information according to SOTER methodology¹⁴. The terrain and terrain component tables were developed from the soil map legend and site characteristics of soil profile cards. In the digitized soil map, locations of each series were georeferenced (Figures 2 and 3) and this map was used to link the attribute information to the soil map unit.

Some of the mandatory information required by SOTER such as bulk density (BD), moisture content at different tensions and saturated hydraulic conductivity (sHC) were not available in some of the published soil survey reports from which the series information was

extracted. Therefore, pedo-transfer functions (PTFs) were developed²⁷ for water retention characteristics at different suctions, BD and sHC, in order to generate this information. Step-wise multiple regression technique in a statistical software (SPSS: version 18.0) was used to develop multiple regression models as PTFs. Scatter-plot diagrams were used to identify the variables and develop a working hypothesis about their relationships. The independent variables used for PTFs were selected considering the cause-effect relationship and correlation coefficient amongst the variables. Details of the procedures and variables selected are explained elsewhere²⁷. The PTFs were validated separately for both the IGP and BSR with a set of available data on these parameters.

All pedon observations were coded in a format to identify the profile that is to be linked with the map (Figure 4). Thus 144 pedons from the IGP and 101 pedons from the BSR were linked to the map (Tables 4 and 5). These are distributed in 11 AESRs and five bioclimates of the IGP ranging from per-humid to arid through sub-humid and semi-arid. Thus the SOTER IGP and BSR cover 72 and 54% of the area of these regions respectively. In SOTER, there are provisions to upload the information when reliable data are available. However, the profile information collated so far can be utilized for the development of the SOTER database for different states on the 1 : 250,000 scale in future, wherein the map units will be greater in number. SOTER is a comprehensive database, which includes all soil and site characteristics, including climate and vegetation. Climate data for 82 climate stations, which were based on point observation, were compiled in the SOTER format and the link between soil and terrain information was made using geographical coordinates.

Application of SOTER database: An advantage of any database developed from the measured soil properties is that it includes data on which dependable decision relating to the most appropriate uses and management of soils can be made. Information synthesized on soils from a

Table 3. Distribution of point data in states, AERs, AESRs and bioclimatic regions of BSR

States	No. of soil series	AERs	AESRs	Bioclimate*	Source
Andhra Pradesh	57	3, 6, 7, 12, 18	3, 6.2, 7.1, 7.2, 7.3, 12.2, 18.3, 18.4	A, SAm, SAd, SHd	9, 45, 57, 61–65
Assam	2	15	15.2	PH, H, SHm, SHd	66
Bihar	4	9, 13	9.2, 13.1	SHd	58
Chhattisgarh	14	11, 12	11, 12.1	SHd, SHm	57, 67
Gujarat	33	2, 4, 5, 19	2.4, 4.2, 5.1, 5.2, 19.1	A, SAm, SAd, SHm	9, 45, 63, 65, 68
Karnataka	13	3, 6, 7	3, 6.1, 6.2, 6.4, 7.1, 7.2	SAd	9, 45, 63, 65, 69
Kerala	3	19	19.3	PH,H	70
Madhya Pradesh	86	4, 5, 10, 11, 12	4.3, 4.4, 5.2, 10.1, 10.2, 10.3, 10.4, 11.0, 12.2	SAm, SHd, SHm	9, 45, 63, 65, 71–73
Maharashtra	204	6, 10, 12, 19	6.1, 6.2, 6.3, 6.4, 10.2, 10.4, 12.1, 19.1	SAm, SAd, SHd, H	9, 63, 65, 73–88
Odisha	6	12, 18	12.1, 12.2, 18.4, 18.5	SHm	89
Punjab	1	2, 9	2.3	SAd, SAm, A	45
Rajasthan	14	2, 4, 5	2.1, 4.2, 5.2	SAd, SAm	9, 63, 90
Tamil Nadu	7	8, 18	8.1, 8.2, 8.3, 18.2	SAm	9, 57, 63, 65
Uttar Pradesh	1	4	4.3	SHm, SHd, SAm, SAd	45
West Bengal	3	15	15.1	SHm	45, 59
Jammu & Kashmir	–	14		A, SAm, SHd, SHm	
Total	448 (425)**		36		

*H, Humid; SHm, Sub-humid moist; SHd, Sub-humid dry; SAm, Semi-arid moist; SAd, Semi-arid dry, A, arid.

**No. of points georeferenced. (many points overlapped due to scale limitation)




Figure 4. Keys for identification of profile from the master datasets in SOTER.




Figure 5. Soil bulk density (0–30 cm) map developed for the Indo-Gangetic Plains using SOTER datasets.

map or stored in an information system, such as SOTER, can be used to make decisions for optimum utilization, planning and management of soil and land resources.




Figure 6. Soil bulk density (0–30 cm) map developed for black soil region using SOTER datasets.

Using the base map of IGP and BSR and the corresponding attribute database, a number of thematic maps were prepared and a few of them are explained below.

Bulk density: BD is an important soil physical parameter which regulates the movement of water, air and roots into the soil and it depends on the texture and organic matter content of the soil. Management interventions, particularly cultural practices, have an impact on this

Georeferenced SIS for agricultural LUP

Table 4. Details of soil series selected from the master soil database to develop the SOTER IGP

State	No. of soil series	AESRs covered	Bioclimate*	Soil series
Punjab	20	9.1, 4.1, 2.3	SAd (8) SAm (7) A (5)	Balewal, Langrian, Phagunwala, Tulewal, Phaguwala, Kanjli, Tulewal, Patiala Dhoda, Nabha, Gurudaspur, Fattu, Mund, Sadhu, Jagjitpur Dewan-Khera, Nihal-Khera1, Nihal-Khera2, Panjgram Kalan, Fatehpur
Haryana	5	4.1, 9.1, 2.3	SAd (2) SAm (2) A (1)	Lukhi, Zarifa Viran Shahazadpur, Berpura Hisar
Uttarakhand	1	9.2	SAm (1)	Haldi
Delhi	3	4.1	SAd (3)	Daryapur, Kakra, Holambi
Uttar Pradesh	33	9.1, 4.1, 4.3, 9.2, 13.1	SAd (2) SAm (26) SHd (5)	Shergarh, Garhsauli Shekhupur, Jauli, Mawana, Hirapur, Nandpur, Maktaura, Charpur, Allahpur, Tilhar, Saunda, Garcia, Bikranpur, Kabirpur, Gopalpur, Lajjanagar, Makhanpur, Nangla Bhagat, Sehud, Ajlapur, Chamkani, Nagaria, Bijaipur, Nagla Jola, Masuri, Parichhatgarh, Khanpur Haderpur, Zikhripur, Jangipur, Tikarikhurd, Basiaram
Bihar	45	13.1, 9.2	SHm (15) SHd (30)	Darwabari, Dumri, Fatehpur, Bananiya, Bhawanipur, Belgachhi, Ganeshpur, Haldikhora, Karamwa, Khanua, Arraha, Bhargaon, Madhuban, Nirpur, Tikapatti Bikramganj, Dullahpur, Baruna, Budhauri, Dadar, Datraul, korap, Dahiya, Katharain, Pokhrahi, Shivpur, Sarthua, Mathiya, Baswariya, Gandhrain, Ghoga, Qutubpur, Sagauli, Sangrampur, Walipur, Bahera, Parsouna, Pipra Naurangia, Narayanpur, Tarapur, Baratol, Hirapatti, Gaupur, Nanpur, Ekchhari
West Bengal	35	12.3, 15.1, 16.2, 15.3, 13.1	SHm (27) PH (8)	Majiara, Silampur, Balidanga, Ghoshat, Balia, Nampur, Alinagar, Salmara, Samaspur, Gangarampur, Barabarua, Sahazadpur, Tulsidanga, Ruisanda, Gopalpur, Balrampur, Chakprayag, Panchpota, Harinathpur, Jatikrishnapur, Jambani, Kanaidighi, Belar, Madhupur, Madhpur, Amarapur, Bansghata Singvita, Chunabhati, Daraboyjot, Berubari, Binnaguri, Matiarkuthi, Seoraguri, Mechpara
Tripura	2	17.2	PH (2)	Nayanpur, Khowai

*PH, Per-humid; SHm, Sub-humid moist; SHd, Sub-humid dry; SAm, Semi-arid moist; SAd, Semi-arid dry; A, arid; number of soil series is shown in parentheses.

property. In rice and potato-growing regions of West Bengal, an increase in BD immediately below the plough layer was observed, which has adversely affected the potato yield²⁸. Similar observations were also reported in rice and wheat-growing soils of Punjab²⁹, some cotton-growing soils of Maharashtra (where subsoil sodicity is prevalent) and the soybean-wheat growing Vertisols of Madhya Pradesh³⁰. Theme maps (Figures 5 and 6) for BD at different depths were developed for both the IGP and the BSR using SOTER. BD at the surface (0–30 cm depth) in soils of the IGP is $> 1.6 \text{ Mg m}^{-3}$ in 81% of the area, wherein the organic carbon is $< 0.50\%$. This, low organic carbon is correlated with high BD, which adversely affects crop growth. High BD is the result of mechanized farming and the use of water with high salt content for many years. BD decreases with depth. The mechanized and intensive agriculture in these areas may form hard pans at the surface or subsurface layers and make the soil impervious to water, air and roots. In contrast, in the BSR, the occurrence of soils with BD

$> 1.6 \text{ Mg m}^{-3}$ is negligible, indicative of low rate of adoption of intensive management practices in these areas.

Bhattacharyya *et al.*³⁰ reported an inverse relationship of BD with soil organic carbon (SOC) in soils of the semi-arid tropics, which increased in soils of drier bioclimates in accordance with the increase in soil inorganic carbon and subsoil sodicity. Therefore, proper rehabilitation measures are required to control BD, porosity and sHC, particularly in the rainfed areas.

Saturated hydraulic conductivity: The hydraulic properties of soils have important implications for management as they regulate water–air relationships and also nutrient availability. The soils of the IGP are well known for their rice–wheat cropping systems and recent reports indicate that production has plateaued or declined due to the emergence of some soil-related constraints. The rice crop of IGP requires standing water, unlike wheat. In contrast to this, cropping in major parts of BSR depends on the water stored in the profile during the monsoon and its

Table 5. Details of soil profiles (series) selected from the master soil database for development of the SOTER BSR

State	No. of soil series	AESRs covered	Bioclimate*	Soil series
Andhra Pradesh	10	3, 6.2, 7.2, 18.4	A (2) SAm (6)	Sollapuram, Tatireddipalle Hugaluru, Amalapuram, Bhimavaram, Jammalamadugu, Pangidi, Nipani
Bihar	4	9.2, 13.1	SAd (1) SHd (1)	Jajapur 1 Mummadivaram
Chhattisgarh	7	11, 12.1	SHd (4) SHm (3)	Barew, Bhadasi, Belsar, Dahiya Pendri Kalan, Hirawani, Boda, Bichanpur Umariguda, Sirgeri, Khujji Kalam
Gujarat	12	2.4, 4.2, 5.1, 5.2, 19.2	A (2) SAm (2) SHd (7) SHm (1)	Bhimdevka, Sokhada 1 Haladar, Mulad Ratanvav, Arnej, Bhola, Chabhadia, Dholi, Jalia, Kumbhara Eru
Karnataka	3	6.4, 7.1, 7.2	SAd (3)	Teligi, Achmatti, Raichur
Madhya Pradesh	33	4.4, 5.2, 10.1, 10.3, 10.4, 11.0, 12.2	SAm (11) SHm (9) SHd (13)	Loni, Bijapur Kalan, Gopalpur, Sarol, Namali, Shankarali, Baiharai, Digwar, Deorikalam, Bainar, Arsia Gonditola, Tejgarh, Kheri, Gaintara, Semarar, Chhapratola, Chandranagar, Karloka, Makajhir Savli, Nabibagh, Talgaon, Baroda Kalam, Mariyadar, Padariya, Madanpur, Sumariyakalam, Jamra, Amziri, Sundra, Rohana 2, Lalatora-1
Maharashtra	15	6.1, 6.2, 6.3, 10.2, 10.4, 12.1, 19.1	SAm (2) SAd (5) SHd (7) H (1)	Anjana, Bhugaon Satgaon, Dhulgaon, Purandarwada, Masala, Nimone, Boripani, Jamnapur, Bhis, Sindewahi, Andhali, Bahadura, Lasanpur Palghar
Odisha	4	12.1, 18.4, 18.5	SHm (4)	Birsinghasahi, Sanfansi, Daiapalli, Nalibasant
Rajasthan	9	4.1, 5.2	SAd (1) SAm (8)	Datwasa Sunel Chhoti, Khando, Raipur, Khanpur, Anta, Kushalgarh, Arnod, Jalawara
Tamil Nadu	2	8.1	SAm (2)	Kovilpatti, Salur
West Bengal	2	15.1	SHm (2)	Hanrgram, Gopalpur

*H, humid; SHm, Sub-humid moist; SHd, Sub-humid dry; SAm, Semi-arid moist; SAd, Semi-arid dry; A, Arid. Number of soil series is shown in parentheses.




Figure 7. Saturated hydraulic conductivity (sHC) map for the Indo-Gangetic Plains.

release during the crop growth. Kadu *et al.*³¹ observed that sHC can be considered as a robust parameter for the determination of the suitability of deep black soils for rainfed cotton and they reported a significant reduction in

yield when sHC decreases to $<10 \text{ mm h}^{-1}$. Therefore, theme maps for sHC are important from an agricultural crop growth point of view in both the IGP and BSR.

sHC (0-100 cm depth) in IGP soils is low ($<2 \text{ mm h}^{-1}$) in major areas (45%; Figure 7), indicating poor drainage to favour rice crop which requires standing water. However, low sHC may pose a threat due to soil and water erosion and flooding during the monsoon and a low storage of soil water for post-rainy season.

In the BSR, the sHC map is presented for the entire 100 cm depth because the major part of the area is under rainfed conditions (Figure 8). Considering sHC of 10 mm h^{-1} as threshold for good agricultural land, an area of 20% that has $<10 \text{ mm h}^{-1}$, needs immediate management interventions to improve drainage. Some BSR areas have pedogenic CaCO_3 and subsoil sodicity^{11,32}. The sub-soil sodicity causes dispersion of clay (which is very high in Vertisols) and clogs the micropores and subsequently decreases hydraulic properties. However, the constant release of Ca^{2+} ions from zeolites and gypsum improves the sHC and overshadows the ill effects of sodium and

Georeferenced SIS for agricultural LUP

magnesium^{11,13} in the BSR under semi-arid and arid climates.

Soil organic carbon: The status of organic carbon in soil is considered as an indicator of soil quality in tropical soils because of its influence on soil properties and crop production³³. Surface soils of IGP and BSR (Figures 9 and 10) indicate >95% soils have organic carbon of <1%, which is considered as a threshold value for a sufficient level in soils of tropical India³⁴. Therefore, soils of IGP and BSR need immediate management intervention, particularly in those areas where organic carbon is <0.5% (85% of area in the IGP and 38% of the area in BSR).




Figure 8. Saturated hydraulic conductivity map for the black soil region.




Figure 9. Soil organic carbon (0–30 cm) status map of the Indo-Gangetic Plains.

The climatic aridity and subsequent development of pedogenic carbonate in the major part of IGP, along with intensive agriculture with low organic inputs are the probable reason for their low organic carbon status. However, studies with limited datasets and carbon prediction models^{23,35} indicate that there is a trend of increase in organic carbon stock in the soils of the IGP and BSR, which followed an initial decline during the post-green revolution period. Reports also indicate that soils of the arid and semi-arid climate, occupying more than one-third of the area of IGP, are prone to be calcareous and sodic due to low levels of organic carbon. Proper rehabilitation programmes could make these soils resilient and thus improve their quality through carbon sequestration^{11,23,36}. Vast areas of arid, semi-arid and sub-humid bioclimates of BSR are low in organic carbon, but have a high potential to sequester more organic carbon due to better substrate quality and thus can be prioritized for the sequestration of organic carbon. The thematic map developed through SOTER can be taken as a guide for selection of areas for the development of treatment plans on a priority basis.

Other applications: GeoSIS and the SOTER database can be used for many applications. Terrain parameters are most commonly used as extensively mapped secondary and auxiliary variable to improve spatial prediction of soil-scapes and soil physical properties like thickness of horizon and physical properties^{37,38}. Soil information systems can be combined with digital elevation models and satellite radiometric data for regional soil mapping³⁹.




Figure 10. Soil organic carbon (0–30 cm) status of the black soil region.

The database has been successfully used to refine the AESR map of India⁴⁰, which will help to determine and demarcate the crop efficiency zones. The digital soil resource database along with climate and plant requirement can be combined to evaluate and categorize the land for different uses. The georeferenced database has been used to evaluate the soils of BSR and IGP for cotton and soybean systems and rice–wheat system respectively⁴¹ in the benchmark spots. The database along with ancillary datasets are also used for predicting the yield of cotton and rice in BSR and the IGP soils respectively⁴². It will also help in assessing the soil and evolving land quality parameters and strategies to improve the quality and health of soils for better use on a sustainable basis⁴³. The primary database has also been widely used to develop PTFs for estimating the physical properties like BD and SHC of soils²⁷, which are seldom available in routine soil survey reports. The other applications of SOTER include assessment of soil erosion, land degradation studies, conservation strategies, land productivity potential, spatial decision support system and environment protection studies.

Conclusions

A GeoSIS in the SOTER framework developed for the IGP and BSR forms a robust database which includes morphological, physical and chemical characteristics of soils along with the site characteristics and climate related information. The information from this organized database can be easily retrieved for use and is compatible with the global database. Although each map unit in the SOTER database can be linked with one profile information, the information collected can be utilized when we develop the SOTER for individual states on 1 : 2,50,000 scale. This warehouse of organized soil and land resource information would form the basis for the development of a SOTER database for the entire country. This can be updated with the recent and relevant data. It is expected that this robust database in a structured framework can be utilized by many users for future scientific and resource planning purposes and thus can remain as a national and international reference database.

1. Nachtergaele, F. and van Ranst, E., Qualitative and quantitative aspects of soil databases in tropical countries. In *Evolution of Tropical Soil Science: Past and Future* (ed. Stoops, G.). Koninklijke Academie voor oversee Wetenschappen, Brussel, 2002, pp. 107–126.
2. Lagacherie, P. and McBratney, A. B., Spatial soil information system and spatial soil inference systems: perspectives for digital soil mapping. *Digital Soil Mapping 2004 Montpellier*, 13–17 September 2004, Elsevier, 2005.
3. Purnell, M. F., Soil survey information supply and demand: international policies and stimulation programmes. *ITC J.*, 1993, **1**, 30–35.
4. Zinck, A. J., Introduction. Special Issue on Soil Survey Workshop, *ITC J.*, 1993, **1**, 2–7.
5. Winegardner, D. L., *An Introduction to Soils for Environmental Professions*, CRC Press, Boca Raton, 1995.
6. Hartemink, A. E., McBratney, A. B. and Mendonca-Santos, M. L. (eds), *Digital Soil Mapping with Limited Data*, Springer, Berlin, 2008.
7. McBratney, A. B., Minasny, B., Cattle, S. R. and Vervoort, R. W., From pedotransfer functions to soil inference systems. *Geoderma*, 2002, **109**, 41–73.
8. Bhattacharyya, T. and Mandal, B., Soil information system of the Indo-Gangetic Plains for resource management. ISSS special session on land use planning. *J. Indian Soc. Soil Sci., Platinum Jubilee Symposium-Proceedings*, 2009, 1–19.
9. Pal, D. K., Bhattacharyya, T. and Bhuse, S. K., Developing a model on the formation and resilience of naturally degraded black soil of Peninsular India as a decision support system for better land use planning. National Bureau of Soil Survey and Land Use Planning, Nagpur and DST, Government of India, Final report to DST, 2003, p. 144.
10. Kolhe, A. H., Chandran, P., Ray, S. K., Bhattacharya, T., Pal, D. K. and Sarkar, D., Genesis of associated red and black shrink – swell soils of Maharashtra. *Clay Res.*, 2011, **30**, 1–11.
11. Pal, D. K., Bhattacharyya, T., Ray, S. K., Chandran, P., Srivastava, P., Durge, S. L. and Bhuse, S. R., Significance of soil modifiers (Ca-zeolites and gypsum) in naturally degraded Vertisols of Peninsular India in identifying the sodic soils. *Geoderma*, 2006, **136**, 210–228.
12. Bhattacharyya, T., Pal, D. K., Srivastava, P. and Velayutham, M., Natural zeolites as savior against soil degradation. *Gondwana Geol. Sci. Mag.*, 2000, **16**, 27–29.
13. Bhattacharyya, T., Pal, D. K., Lal, S., Chandran, P. and Ray, S. K., Formation and persistence of Mollisols on zeolitic Deccan basalt of humid tropical India. *Geoderma*, 2006, **136**, 609–620.
14. Van Engelen, V. W. P. and Wan, T. T. (eds), *Global and National Terrain Digital Database (SOTER) Procedure Manual* (revised edition), International Soil Reference and Information Centre, Wageningen, The Netherlands, 1995.
15. Batjes, N. H. and Dijkshoorn, I. A., Carbon and nitrogen blocks in the soils of the Amazon region. *Geoderma*, 1999, **89**, 273–286.
16. Falloon, P. D., Smith, P., Smith, J. V., Szabo, J., Coleman, K. and Marshall, S., Regional estimates of C sequestration potential: linking the Rothamsted C model to GIS databases. *Biol. Fert. Soils*, 1998, **27**, 236–241.
17. Batjes, N. H. *et al.*, Preparation of consistent soil data sets for modeling purposes: secondary SOTER data for four case study areas. In *Soil Carbon Stocks at Regional Scales* (eds Milne, E., Powlson, D. S. and Cerri, C. E. P.), *Agric. Ecol. Environ.*, 2007, vol. 122, pp. 26–34.
18. Chandran, P. *et al.*, SOTER (Global and national soil and terrain digital databases) IGP, India. Special publication for ‘Assessment of Soil Organic Carbon Stock and Change at National Scale’, NBSS&LUP, India, 2005, p. 101.
19. Staff, NBSS&LUP, Soils of India, NBSS Publ. No. 94, National Bureau of Soil Survey & Land Use Planning, Nagpur, 2002, pp. 130 + 11 sheet maps.
20. Mandal, *et al.*, Revisiting agro-ecological sub-regions of India – a case study of two major food production zones. *Curr. Sci.*, 2014, **107**(9), 1519–1536.
21. Sehgal, J., Mandal, D. K., Mandal, C. and Vadivelu, S., Agro-ecological regions of India, NBSS&LUP, Nagpur, Tech. Bull. Publ., 24, 1992, 2nd edn, p. 130.
22. Velayutham, M., Mandal, D. K., Mandal, C. and Sehgal, J. L., Agro-ecological subregion of India for development and planning, NBSS Publ. No. 235, National Bureau of Soil Survey and Land Use Planning, Nagpur, 1999, p. 450.
23. Bhattacharyya, T., Pal, D. K., Chandran, P., Mandal, C., Ray, S. K., Gupta, R. K. and Gajbhiye, K. S., Managing soil carbon stock

- in the Indo-Gangetic Plains, India. Rice–Wheat Consortium for the Indo-Gangetic Plains, New Delhi, 2004, p. 44.
24. Ray, S. K. *et al.*, Baseline data Indo-Gangetic Plains (IGP) project ‘Georeferenced soil information system for land use planning and monitoring soil and land quality for agriculture’. Working report No. 1, Parts I & II, NAIP Component-4, NBSS&LUP, Nagpur, 2014.
 25. Bhattacharyya, T. *et al.*, Soil datasets of the hot spots Indo-Gangetic Plain (IGP) project ‘Georeferenced soil information system for land use planning and monitoring soil and land quality for agriculture’, Working report No. 3, NAIP component-4, NBSS&LUP, Nagpur, 2014.
 26. Bhattacharyya, T. *et al.*, Soils of India: historical perspective, classification and recent advances. *Curr. Sci.*, 2013, **104**, 1308–1323.
 27. Tiwary, P. *et al.*, Pedotransfer functions: a tool for estimating hydraulic properties of two major soil types of India. *Curr. Sci.*, 2014, **107**(9), 1431–1439.
 28. Bhattacharyya, T., Ray, S. K., Sahoo, A. K., Durge, S. L., Chandran, P., Sarkar, D. and Pal, D. K., Pan formation in soils under paddy–potato/mustard–paddy systems in Indo-Gangetic Plain. West Bengal Rice–Wheat Consortium for the Indo-Gangetic Plains, 2006; www.rwc.cgair.org/Pub_Info.asp?ID=165
 29. Sidhu, G. S. *et al.*, Impact of management levels and land-use changes on soil properties in rice–wheat cropping system of the Indo-Gangetic Plains. *Curr. Sci.*, 2014, **107**(9), 1487–1501.
 30. Bhattacharyya, T. *et al.*, Carbon sequestration in red and black soils. II. Influence of physical and chemical properties. *Agrope-dology*, 2007, **17**, 16–25.
 31. Kadu, P. R., Vaidya, P. H., Balpande, S. S., Satyavathi, P. L. A. and Pal, D. K., Use of hydraulic conductivity to evaluate the suitability of Vertisols for deep rooted crops in semi-arid parts of central India. *Soil Use Manage.*, 2003, **19**, 208–216.
 32. Pal, D. K., Dasog, G. S., Vadivelu, S., Ahuja, R. L. and Bhattacharyya, T., Secondary calcium carbonate in soils of arid and semi-arid regions of India. In *Global Climate Change and Pedogenic Carbonates* (eds Lal, R. *et al.*), Lewis Publishers, Boca Raton, FL, USA, 2000, pp. 149–185.
 33. Chandran, P., Ray, S. K., Durge, S. L., Raja, P., Nimkar, A. M., Bhattacharyya, T. and Pal, D. K., Scope of horticultural land use system in enhancing C sequestration in ferruginous soils of semi-arid tropics. *Curr. Sci.*, 2009, **97**, 1039–1046.
 34. Velayutham, M., Pal, D. K. and Bhattacharyya, T., Organic carbon stock in soils of India. In *Global Climatic Change and Tropical Ecosystems* (eds Lal, R., Kimble, J. M. and Stewart, B. A.), Lewis publishers, Boca Raton, FL, USA, 2000, pp. 71–96.
 35. Bhattacharyya, T., Chandran, P., Ray, S. K., Pal, D. K., Venugopalan, M. V., Mandal, C. and Wani, S. P., Changes in levels of carbon in soils over years of two important production zones of India. *Curr. Sci.*, 2007, **93**, 1854–1863.
 36. Gupta, R. K. and Rao, D. L. N., Potential of wastelands for sequestering carbon by representation. *Curr. Sci.*, 1994, **66**, 378–380.
 37. Moore, I. D., Gessler, P. E., Nielsen, G. A. and Peterson, G. A., Soil attribute prediction using terrain analysis. *Soil Sci. Soc. Am. J.*, 1993, **57**, 443–452.
 38. Gessler, P. E., Moore, I. D., McKenzie, N. J. and Ryan, P. J., Soil landscape modeling and spatial prediction of soil attributes. *Int. J. Geogr. Inf. Syst.*, 1995, **9**, 421–432.
 39. Dobos, E., Micheli, E., Baumgardner, M. F., Biehl, L. and Helt, T., Use of combined digital elevation model and satellite radiometric data for regional soil mapping. *Geoderma*, 2000, **97**, 3–4.
 40. Mandal, C. *et al.*, Revisiting agro-ecological subregions of India – a case study of two major food production zones. *Curr. Sci.*, 2014, **107**(9), 1519–1536.
 41. Chatterji, S. *et al.*, Land evaluation for major crops in the Indo-Gangetic Plains and black soil regions using fuzzy model. *Curr. Sci.*, 2014, **107**(9), 1502–1511.
 42. Venugopalan, M. V. *et al.*, InfoCrop-cotton simulation model – its application in land quality assessment for cotton cultivation. *Curr. Sci.*, 2014, **107**(9), 1512–1518.
 43. Ray, S. K. *et al.*, Soil and land quality of the Indo-Gangetic Plains of India. *Curr. Sci.*, 2014, **107**(9), 1470–1486.
 44. Sehgal, J. L. and Sharma, P. K., Benchmark soils of Punjab. Soils Bulletin 5, Department of Soils, Punjab Agricultural University, Ludhiana, 1982, p. 211.
 45. Lal, S., Deshpande, S. B. and Sehgal, J. (eds), Soil Series of India. National Bureau of Soil Survey and Land Use Planning, Nagpur, India. NBSS Publ 40, 1994, p. 684.
 46. Ray, S. K. *et al.*, Benchmark soil series of the Indo-Gangetic Plains (IGP), India. Special publication for ‘Assessment of soil organic carbon stock and change at national scale’. NBSS&LUP, India, 2005, p. 186.
 47. Pal, D. K. *et al.*, Pedogenic thresholds in benchmark soils under rice–wheat cropping system in a climosequence of the Indo-Gangetic Alluvial Plains. Final project report, Division of Soil Resource Studies, NBSS&LUP, Nagpur, 2010, p. 193.
 48. Martin, D., Sharma, J. P., Singh, S. P. and Sarkar, D., Soils resource mapping of Shahjahanpur district of Uttar Pradesh for perspective land use planning. NBSS Publ. No. 428, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2009.
 49. Murthy, R. S., Hirekerur, L. R., Deshpande, S. B., Venkata Rao, B. V. and Shankaranarayana, H. S., Benchmark soils of India – morphology, characteristics and classification for resource management. National Bureau of Soil Survey and Land Use Planning, Nagpur, 1982, p. 374.
 50. Verma, T. P., Singh, S. P., Ram Gopal, Dhankar, R. P., Jagat Ram and Sarkar, D., Soil resource for land use planning Etawah District (UP). National Bureau of Soil Survey and Land Use Planning, Nagpur, NBSS Publ. 2005, p. 124.
 51. Singh, S. K. and Agrawal, H. P., Morphology and taxonomy of prominent soils developed in alluvial pedogenic complex of Gomti-Sai interflue, Dist. Jaunpur, UP. *J. Sci. Res.*, 1995, **45**, 49–61.
 52. Bhargava, G. P., Pal, D. K., Kapoor, B. S. and Goswami, S. C., Characteristics and genesis of some sodic soils in the Indo-Gangetic Plains of Haryana and Uttar Pradesh. *J. Indian Soc. Soil Sci.*, 1981, **29**, 61–70.
 53. Mahapatra, S. K., Sharma, J. P. and Sarkar, D., Soil resource mapping of Mathura district of Uttar Pradesh for perspective land use planning. NBSS Publ., NBSS&LUP, Nagpur, 2010, p. 156.
 54. Jagat Ram, Ram Gopal, Obi Reddy, G. P. and Sarkar, D., Soil resource mapping for land use planning of Moradabad District (UP). National Bureau of Soil Survey and Land Use Planning, Nagpur, NBSS Publ., 2009, p. 159.
 55. Verma, T. P., Singh, S. P., Martin, D., Dhankar, R. P., Sharma, J. P. and Sarkar, D., Soil resource mapping of Mainpuri district of Uttar Pradesh for perspective land use planning. National Bureau of Soil Survey and Land Use Planning, Nagpur, 2009.
 56. Walia, C. S., Sharma, J. P. and Sarkar, D., Soil resource mapping of Firozabad district of Uttar Pradesh for perspective land use planning. National Bureau of Soil Survey and Land Use Planning, Nagpur, 2009, p. 107.
 57. Staff, NBSS&LUP, National register of soil series. National Bureau of Soil Survey and Land Use Planning, Nagpur (unpublished), 2012.
 58. Sahoo, A. K., Sarkar, D. and Gajbhiye, K. S., Soils series of Bihar. Publ. No. 98, National Bureau of Soil Survey and Land Use Planning, Nagpur, NBSS, 2002, p. 289.
 59. Nayak, D. C., Sarkar, D. and Velayutham, M., Soil series of West Bengal, NBSS Publ. No. 89, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2001, p. 260.
 60. Sarkar, D., Nayak, D. C., Maurya, U. K. and Gajbhiye, K. S., Mineralogy of benchmark soils of West Bengal, Publ. No. 139, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, NBSS, 2008, p. 171.

61. Reddy, R. S., Budihal, S. L., Ramesh Kumar, S. C. and Naidu, L. G. K., Benchmark soils of Andhra Pradesh, NBSS Publ no. 128, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2005, p. 143.
62. Naidu, L. G. K., Hegde, R., Ramamurthy, V., Thayalan, S., Srinivas, S. and Niranjana, K. V., Soil resources of Anantapur District, Andhra Pradesh, NBSS Publ. No. 1017, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2008, p. 22.
63. Bhattacharyya, T., Pal, D. K., Chandran, P., Ray, S. K., Mandal, C. and Telpande, B., Soil carbon storage capacity as a tool to prioritize areas for carbon sequestration. *Curr. Sci.*, 2008, **95**, 482–494.
64. Naidu, L. G. K., Niranjana, R. V., Ramamurthy, V., Ramesh Kumar, S. C., Anilkumar, K. S. and Thayalan, S., Planning optimum land use based on biophysical and economic resource of Pulivendala Agricultural Div., Kadapa Dist, Andhra Pradesh, NBSS Publ. No. 1029, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2009, p. 184.
65. Bhattacharyya, T. *et al.*, Soil datasets of the hot spots black soil region (BSR) project on 'Georeferenced soil information system for land use planning and monitoring soil and land quality for agriculture'. NAIP Component 4, Working report no. 4, NBSS&LUP, Nagpur, 2014, p. 228.
66. Vadivelu, S., Sen, T. K., Bhaskar, B. P., Baruah, U., Sarkar, D. and Maji, A. K., Soil series of Assam, NBSS Publ. No. 101, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2004, p. 229.
67. Tamgadge, D. B., Gajbhiye, K. S. and Pande, G. P., Soils series of Chhattisgarh, NBSS Publ. No. 85, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2002, p. 183.
68. Sharma, J. P., Giri, J. D., Shampura, R. L. and Gajbhiye, K. S., Soil series of Gujarat, NBSS publ no. 120, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2006, p. 399.
69. NBSS&LUP Staff, Soils of Bellary District, NBSS Publ. No. 1015, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2008, p. 172.
70. Nair, K. M. *et al.*, Soils of Elapully panchayat, Palakkad, Kerala, NBSS Publ. No. 962, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2006, p. 58 + Annexure.
71. NBSS&LUP Staff, Soils of Palassdoh Minor, Rohana Minor 2 and 3 in Etawa Command, Hoshangabad, Madhya Pradesh, NBSS Publ. No. 532, National Bureau of Soil Survey and Land Use Planning, Nagpur, 1996, p. 80.
72. Tamgadge, D. B., Gajbhiye, K. S., Velayutham, M. and Kaushal, G. S., Soil series of Madhya Pradesh, NBSS Publ. No. 78, National Bureau of Soil Survey and Land Use Planning, Nagpur, 1999, p. 523.
73. NBSS&LUP Staff, Soils of Lalatora watershed, District Vidisha, Madhya Pradesh, NBSS Publ. No. 549, National Bureau of Soil Survey and Land Use Planning, Nagpur, 1999, p. 60.
74. NBSS&LUP Staff, Soils of Bhandra District, Maharashtra, NBSS Publ. No. 520, National Bureau of Soil Survey and Land Use Planning, Nagpur, 1991, p. 33.
75. Kadu, P. R., Physical and chemical properties affecting hydraulic conductivity of black soils of the Purna valley. MSc thesis, Dr Panjabrao Deshmukh Krishi Vidhyapeeth, Akola, 1991, p. 54.
76. Pacharne, T. K., Characteristics and genesis of associated red and black soils of the part of the Saptdhara watershed. M Sc thesis, Dr Panjabrao Deshmukh Krishi Vidhyapeeth, Akola, 1992, p. 53.
77. NBSS&LUP Staff, Soils of Chandrapur District, Maharashtra, NBSS Publ. No. 525, National Bureau of Soil Survey and Land Use Planning, Nagpur, 1993, p. 61.
78. Balpande, S. S., Characteristics, genesis and degradation of Vertisols of Purna valley, Maharashtra. Ph D thesis, Dr Panjabrao Deshmukh Krishi Vidhyapeeth, Akola, 1993, p. 193.
79. Paranjape, M. V., Study of waterlogged soils of the World Bank Watershed Project (Wrwbd-5) near Tamasvada, Wardha District. M Sc thesis, Dr Panjabrao Deshmukh Krishi Vidhyapeeth, Akola, 1995, p. 69.
80. Gabhane, V. V., Occurrence of shallow to deep black soils on the basaltic plateau of Malegaon – their genesis and land use potential. Ph D thesis, Dr Panjabrao Deshmukh Krishi Vidhyapeeth, Akola, 1996, p. 189.
81. Challa, O., Gajbhiye, K. S. and Velayutham, M., Soil Series of Maharashtra, NBSS Publ. No. 79, National Bureau of Soil Survey and Land Use Planning, Nagpur, 1999, p. 429.
82. Balpande, H. S., Assessment of land qualities of grape growing soil in Maharashtra. M Sc thesis, Dr Panjabrao Deshmukh Krishi Vidhyapeeth, Akola, 2000, p. 135.
83. Vaidya, P. H., Evaluation of shrink–swell soils and groundwater of the Pedhi water-shed in Amravati district for land use planning. Ph D thesis, Dr Panjabrao Deshmukh Krishi Vidhyapeeth, Akola, 2001, p. 167.
84. Ashok Kumar, H. P., Characterization and classification of sugarcane growing soils of Ahmadnagar District of Maharashtra. M Sc thesis, Dr Panjabrao Deshmukh Krishi Vidhyapeeth, Akola, 2007, p. 82.
85. Zade, S. P., Pedogenic studies of some deep shrink–swell soils of Marathwada region of Maharashtra to develop a viable land use plan. Ph D thesis, Dr Panjabrao Deshmukh Krishi Vidhyapeeth, Akola, 2007, p. 224.
86. Sharma, J. P., Raja, P., Nair, K. M., Bhaskar, B. P. and Sarkar, D., Reconnaissance soil survey mapping correlation and classification of Wardha District, Maharashtra, NBSS Publ. No. 595, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2008, pp. 126 + 1 map.
87. Deshmukh, V., Determination of layer charge after removal of hydroxy-interlayers in some shrink–swell soil clays of Maharashtra. M Sc thesis, Dr Panjabrao Deshmukh Krishi Vidhyapeeth, Akola, 2009, p. 75.
88. Kolhe, A. H., Characteristics and genesis of red swell–shrink soils of Hingoli district of Maharashtra. M Sc thesis, Dr Panjabrao Deshmukh Krishi Vidhyapeeth, Akola, 2010, p. 91.
89. Sarkar, D., Sah, K. D., Sahoo, A. K. and Gajbhiye, K. S., Soil series of Orissa. NBSS Publ. No. 119, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2005, p. 254.
90. Shyampura, R. L., Singh, S. K., Singh, R. S., Jain, B. L. and Gajbhiye, K. S., Soil series of Rajasthan, NBSS Publ. No. 95, National Bureau of Soil Survey and Land Use Planning, Nagpur, 2002, p. 364.

ACKNOWLEDGEMENTS. The present study was carried out by the National Agricultural Innovative Project (NAIP) (Component 4), sponsored research on 'Georeferenced soil information system for land use planning and monitoring soil and land quality for agriculture' through Indian Council of Agricultural Research, New Delhi. The financial assistance is gratefully acknowledged.