



# A fourth principle is required to define Conservation Agriculture in sub-Saharan Africa: The appropriate use of fertilizer to enhance crop productivity



B. Vanlauwe<sup>a,\*</sup>, J. Wendt<sup>b</sup>, K.E. Giller<sup>c</sup>, M. Corbeels<sup>d,e</sup>, B. Gerard<sup>f</sup>, C. Nolte<sup>g</sup>

<sup>a</sup> Natural Resource Management Research Area, International Institute of Tropical Agriculture (IITA), Nairobi, Kenya

<sup>b</sup> East and Southern Africa Division, International Fertilizer Development Center (IFDC), Nairobi, Kenya

<sup>c</sup> Plant Production Systems, Wageningen University, Wageningen, The Netherlands

<sup>d</sup> Annual Cropping Systems, Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD), Montpellier, France

<sup>e</sup> Embrapa-Cerrados, Planaltina, Brazil

<sup>f</sup> Global Conservation Agriculture Program, International Maize and Wheat Improvement Center (CIMMYT), El Batán, Mexico

<sup>g</sup> Plant Production and Protection Division, Food and Agriculture Organization of the United Nations (FAO), Rome, Italy

## ARTICLE INFO

### Article history:

Received 28 September 2013

Received in revised form 6 October 2013

Accepted 6 October 2013

### Keywords:

Mulch  
Integrated Soil Fertility Management  
Smallholder farmers  
Organic resources

## ABSTRACT

Intensification of agricultural systems in sub-Saharan Africa (SSA) is considered a pre-condition for alleviation of rural poverty. Conservation Agriculture (CA) has been promoted to achieve this goal, based on three principles: minimum tillage, soil surface cover, and diversified crop rotations. CA originated in regions where fertilizer is commonly used and crop productivity is high, ensuring an abundance of crop residues. By contrast, crop yields are generally low in SSA and organic residues in short supply and farmers face competing demands for their use. Since minimal tillage without mulch commonly results in depressed yields, the use of fertilizer to enhance crop productivity and organic residue availability is essential for smallholder farmers to engage in CA. This is especially true since alternative ways to increase organic matter availability have largely failed. A case study from Kenya clearly demonstrates how fertilizer increases maize stover productivity above thresholds for minimal initial soil cover required for initiating CA (about 3 tonne ha<sup>-1</sup>). We conclude that strategies for using CA in SSA must integrate a fourth principle – the appropriate use of fertilizer – to increase the likelihood of benefits for smallholder farmers.

© 2013 Elsevier B.V. All rights reserved.

## 1. Introduction

In recent years, Conservation Agriculture (CA) has been promoted to intensify smallholder farming system sustainably in sub-Saharan Africa (SSA) (Benites and Ashburner, 2003; FAO, 2011). Conservation Agriculture is commonly defined around a set of three principles: minimum tillage, soil surface cover, and diversified crop rotations. One of the main justifications for promoting CA in Africa is its widespread use in large-scale farming in various parts of the world (Bolliger et al., 2006; Kassam et al., 2009), with some adoption by smallholders, e.g., in southern Brazil and Paraguay (Evers and Agostini, 2001). This has been partly driven by the presence of an enabling environment, including the availability of herbicides. It is important to note, however, that definitions of 'smallholder farmers' are not consistent. In southern Brazil, a smallholder is classified

as farming less than 50 ha, whereas in SSA, smallholder farmers commonly have access to less than 2 ha. While some of the initially hypothesized benefits of CA including soil carbon sequestration and increased crop yields have not been unequivocally confirmed (Govaerts et al., 2009; Luo et al., 2010), CA often results in more stable and economically favourable yields, usually after a number of years after conversion from conventional agriculture (Knowler and Bradshaw, 2007; Rusinamhodzi et al., 2011).

Uptake of CA by smallholders in SSA remains limited (Kassam et al., 2009; Andersson and D'Souza, 2013) and a number of important constraints to widespread adoption have been highlighted. The lack of organic resources to provide sufficient surface mulch consistently ranks amongst the top constraints (Erenstein, 2002; Giller et al., 2009), especially in areas with high livestock feed requirements (Valbuena et al., 2012). Minimal tillage without surface mulch usually results in depressed yields (Verhulst et al., 2011; Baudron et al., 2012; Thierfelder et al., 2013), partly because mulch provides the necessary conditions for reduced run-off and soil moisture conservation, notably in drier climates (Mupangwa et al., 2012). Mulch also ensures that the physical conditions of

\* Corresponding author at: IITA, c/o ICIPE, PO Box 30772-00100, Nairobi, Kenya. Tel.: +254 723 972250.

E-mail address: [b.vanlauwe@cgiar.org](mailto:b.vanlauwe@cgiar.org) (B. Vanlauwe).

the topsoil are conducive for seed germination and initial crop growth.

In this paper we argue that a fourth principle – the appropriate use of fertilizer – is required to define CA to enhance both crop productivity and produce sufficient crop residues to ensure soil cover under smallholder conditions in SSA. This paper is not meant to advocate CA but argues that without acknowledging this fourth principle the chance that CA will achieve success, especially with smallholder farmers, is limited. Fertilizer application is proposed as a separate principle for CA in contrast to other agronomic practices, including planting time, spacing, and weeding regime, because fertilizer is essential for CA to work, whilst the sub-optimal implementation of other crop management practices do not lead to the failure of CA as such. Moreover, given the reframing of CA in SSA from a resource-saving to a productivity-enhancing paradigm ([www.fao.org](http://www.fao.org), Andersson and D'Souza, 2013), and recognizing that fertilizer use is essential to raise crop productivity on most African soils (e.g., Mwangi, 1997; Abuja Fertilizer Summit, 2006), the proposed fourth principle is fully aligned to this reframing of CA. Although this principle also applies to other continents, we chose to focus this paper on SSA since this is the continent where CA is commonly promoted in areas with little or no availability or use of fertilizer.

## 2. The origin of CA and its three principles

The CA revolution in the tropics started in the 1970s with large-scale farmers in Brazil (Landers, 1999), and spread to other countries in Latin America and certain parts of South Asia (e.g., the Indo-Gangetic basin; Gupta and Sayre, 2007). Minimum or no-tillage with mulching was advocated to reduce the impact of rain on exposed soil and consequent soil erosion losses (Landers, 1999; Roose and Barthes, 2001). An important benefit of minimum or no-tillage was the energy saved by eliminating several tillage operations. Fertilizer use was already a widespread practice in these systems where high crop yields were common. Fertilizer combined with reduced tillage results in increased soil surface biomass to provide sufficient soil cover, especially in areas with segregation of crop and livestock production, as is prevalent on large-scale farms in the areas mentioned above. Agro-chemicals, including fertilizer and herbicides, are readily available and widely used in all regions in which CA has been adopted. Diversified crop rotations are an important element of good agricultural practice, not restricted to CA but also embraced by other approaches to sustainable intensification, including Integrated Soil Fertility Management (ISFM) (Vanlauwe et al., 2010) and agroforestry (e.g., Jama et al., 1998). Crop rotations allow the inclusion of nitrogen-fixing legumes and break pest and disease cycles of crops that are too frequently planted in the same field. Yet rotations do not always result in improved pest control, as some pests can be stimulated, for instance when large amounts of readily decomposable mulch are present (Chikowo et al., 2004).

## 3. Smallholder farming and CA


Smallholder farming conditions in SSA differ from large-scale farming situations in many aspects relevant to implementation of CA. First of all, smallholder crop yields are often poor due to the limited use of agro-inputs and labour (e.g., Tittonell and Giller, 2013), with little crop residue produced as a consequence. Secondly, in many smallholder farming systems in SSA, there are competing demands on available crop residues, especially for livestock feed (Giller et al., 2009; Valbuena et al., 2012). Thirdly, where population densities leads to fallow land being virtually absent, management-induced soil fertility gradients are created due to a

concentration of scarcely available organic resources, either direct or processed, e.g., as manure or compost, in small areas, usually around the homestead (Harris, 2002; Tittonell et al., 2005). This further degrades more remote plots through nutrient mining, a lack of organic matter recycling and erosion due to lack of soil cover. The consequence is that at the planting of a subsequent crop, the cover of crop residues falls below the 30% required to reduce inter-rill soil erosion substantially (Allmaras and Dowdy, 1985), even if all of the crop residues are recycled. Since the relationship between erosion reduction and mulch cover is exponential, more cover results in strong reductions of erosion (Erenstein, 2002). Mulch also reduces weed growth although substantially greater soil cover is required to suppress weeds (Naudin et al., 2011). In Central Kenya, Guto et al. (2011) showed that with stover yields below  $3 \text{ t ha}^{-1}$ , or equivalent grain yields below  $2.5 \text{ t ha}^{-1}$  (assuming a harvest index of 0.45), surface cover is less than 30% at the onset of the season (Fig. 1a). Average maize grain yields in Kenya for the period 2006–2011 were  $1.6 \text{ t ha}^{-1}$  ([www.fao.org](http://www.fao.org)) with an equivalent stover yield of  $1.9 \text{ t ha}^{-1}$ ; well below the above threshold. However, maize grain yields can easily reach  $4 \text{ t ha}^{-1}$  (Vanlauwe et al., 2006) in the fertilized homestead plots, which are usually less than 10% of the farm land (Tittonell et al., 2010). Average maize grain yields in SSA varied between  $1.0$  and  $1.8 \text{ t ha}^{-1}$  for the period 2006–2011, except for the southern African region ( $3.8 \text{ t ha}^{-1}$ ) ([www.fao.org](http://www.fao.org)). Thus crop productivity in African smallholder systems is commonly below the threshold required to provide sufficient mulch to implement CA successfully.

## 4. The quest for organic resources

Before the widespread promotion of CA, many attempts were made to enhance the availability of organic resources in smallholder farms. These were mainly driven by the search for low-input agricultural practices and the widespread belief that fertilizers were beyond the reach of African smallholder farmers. Examples include alley cropping systems (e.g., Kang et al., 1981), integration of herbaceous legumes (e.g., Carsky et al., 2001), improved legume tree fallows and biomass transfer systems (e.g., Sanchez et al., 1997). Over time, it was realized that the interest of smallholder farmers in the above technologies was disappointing, commonly due to poor crop response (Hauser et al., 2006), management requirements beyond farmers' labour availability (Dvorak, 1996) and/or the lack of immediate benefits to farmers (e.g., Sidibé, 2005). Further, fertilization with phosphorus and other nutrients is generally required to stimulate productive growth of nitrogen fixing legumes to provide the mulch needed (Giller and Cadisch, 1995). With the recent upsurge of interest in CA, some of these options are again being considered to provide the organic resources required to engage in CA. Some approaches place CA amongst technologies that can intensify agricultural systems with limited or no use of fertilizer (e.g., Garrity et al., 2010). In this context, The Montpellier Panel (2013) classified conservation farming under the heading 'ecological intensification', a paradigm that promotes intensification with reduced use of fertilizer whilst capitalizing on ecological processes (De Schutter, 2010). Obviously, earlier constraints to adoption of these technologies remain when implemented in the context of CA. Moreover, CA is not a 'green technology' that does not require the use of external inputs, since fertilizer and herbicides are intensively used in areas where CA has taken off, and are major factors in its success. Gowing and Palmer (2008) concluded that 'CA does not overcome constraints on low-external-input systems in sub-Saharan Africa'.

Since the Abuja Fertilizer Summit in 2006, fertilizer use has regained emphasis in the context of agricultural intensification in Africa, with a specific focus on maximizing the use efficiency



**Fig. 1.** Relationship between maize stover yield and soil cover at the start of the cropping seasons in Central Kenya (adapted from Guto et al., 2011) (a), and maize stover yields for different field types in three districts of Western Kenya with and without fertilizer (adapted from Vanlauwe et al., 2006) (b). The error bars in (b) are standard errors of the difference to compare stover yields within site, the left bar for the plots with fertilizer and the right bar for the control plots. The horizontal dashed line represents the 30% soil cover limit required for CA. Photographs in the centre give an impression of the maize plots close to harvest on the different field types in Central Kenya.

and profitability of nutrient inputs ([www.agra.org](http://www.agra.org)). Consequently, one of the major arguments that led to the focus on low input agriculture, namely the lack of affordable and timely access to fertilizer, is losing relevance. Fertilizer, when applied appropriately, has been demonstrated to increase crop yields substantially and profitably under smallholder farming conditions (Vanlauwe et al., 2011). 'Appropriate' refers to application of the right type of fertilizer at the right rate, time, and place (Zingore and Johnston, 2013) and accompanied by the right agronomic practices (e.g., time of planting, crop spacing, weed control). 'Appropriate' also refers to avoiding non-responsive soils or soils on which crops do not respond to the application of standard NPK fertilizer due to other constraints limiting crop growth besides the nutrients contained in the fertilizer (Rusinamhodzi et al., 2013).

### 5. Appropriate fertilizer use as a fourth principle for CA in smallholder systems in Africa

The lack of organic resources is a major constraint to adoption of CA by smallholder farmers. Appropriate use of fertilizer results in substantial increases in crop productivity and in the availability of crop residues, across within-farm soil fertility gradients, and more so on mid-distant and remote fields, which constitute the majority of the fields in smallholder farming systems (e.g., Fig. 1b). These findings suggest that crop production in SSA would greatly benefit from fertilizer addition. Enhanced availability of crop residues could also allow alternative uses of these residues while retaining the minimal requirement for soil cover. Since crop productivity in smallholder systems is generally poor, adding the appropriate use of fertilizer as a fourth principle of CA will ensure that valuable farmers' time—which is a limited resource in 'hoe and cutlass' agriculture – is not wasted on interventions with limited scalability. Currently, CA is widely promoted in farming communities that have little access to or use little fertilizer for a variety of reasons. In Zimbabwe, for example, CA is advocated by non-governmental organizations under a range of humanitarian relief programmes. These programmes target the most resource-poor and vulnerable farmers who receive seeds and fertilizer for free, but have virtually no capacity to purchase fertilizers by themselves (Twomlow et al., 2008). Obviously, strategies for promoting CA would need to change if the fourth principle of fertilizer use is considered, with sustainable input supply chains for fertilizer at affordable prices being necessary for success. In this context, ISFM, which aims to assimilate best practices to maximize the use efficiency of fertilizer, could serve as the initial steps towards the adoption of CA by

smallholder farmers, thus moving CA from a 'competitive' to a 'complementary' approach. Further research should focus on defining appropriate fertilizer practices for CA since fertilizer use efficiency will be affected by the other three principles of CA.

### References

- Abuja Fertilizer Summit, 2006. Abuja Declaration on Fertilizer for African Green Revolution. African Union Special Summit of the Heads of State and Government, Abuja, Nigeria.
- Allmaras, R.R., Dowdy, R.H., 1985. Conservation systems and their adoption in the United States. *Soil Till. Res.* 5, 197–222.
- Andersson, J.A., D'Souza, S., 2013. From adoption claims to understanding farmers and contexts: a literature review of Conservation Agriculture (CA) adoption among smallholder farmers in southern Africa. *Agric. Ecosyst. Environ.* (in press).
- Baudron, F., Titttonell, P., Corbeels, M., Letourmy, P., Giller, K.E., 2012. Comparative performance of conservation agriculture and current smallholder farming practices in semi-arid Zimbabwe. *Field Crops Res.* 132, 117–128.
- Benites, J.R., Ashburner, J.E., 2003. FAO's role in promoting conservation agriculture. In: Garcia-Torres, L., Benites, J., Martinez-Vilela, A., Holgado-Cabrera, A. (Eds.), *Conservation Agriculture: Environment, Farmers Experiences, Innovations, Socio-Economy, Policy*. Food and Agriculture Organization of the United Nations, Rome, Italy, pp. 139–153.
- Bolliger, A., Magid, J., Amado, T.J.C., Neto, F.S., Ribeiro, M.D.D., Calegari, A., Ralisch, R., de Neergaard, A., 2006. Taking stock of the Brazilian zero-till revolution: a review of landmark research and farmers' practice. *Adv. Agron.* 91, 47–110.
- Carsky, R.J., Becker, M., Hauser, S., 2001. Mucuna cover crop fallow systems: potential and limitations. In: Tian, G., Ishida, F., Keatinge, J.D.H. (Eds.), *Sustaining Soil Fertility in West-Africa*. SSSA Special Publication Number 58, Madison, USA, pp. 111–136.
- Chikowo, R., Mapfumo, P., Nyamugafata, P., Giller, K.E., 2004. Woody legume fallow productivity, biological N<sub>2</sub>-fixation and residual benefits to two successive maize crops in Zimbabwe. *Plant Soil* 262, 303–315.
- De Schutter, O., 2010. Agro-ecology and the right to food. In: UN Report A/HRC/16/49. United Nations Organization, New York.
- Dvorak, K.A., 1996. Adoption Potential of Alley Cropping. RCMD Research Monograph No.23. International Institute of Tropical Agriculture, Ibadan, Nigeria.
- Erenstein, O., 2002. Crop residue mulching in tropical and semi-tropical countries: an evaluation of residue availability and other technological implications. *Soil Till. Res.* 67, 115–133.
- Evers, G., Agostini, A., 2001. No-Tillage Farming for Sustainable Land Management: Lessons from the 2000 Brazil Study Tour. FAO Investment Centre Occasional Paper Series No. 12. Food and Agriculture Organization of the United Nations, Rome, Italy.
- FAO, 2011. *Save & Grow. A Policymakers Guide to the Sustainable Intensification of Smallholder Crop Production*. Food and Agriculture Organization of the United Nations, Rome, Italy, pp. 116.
- Garrity, D., Akinnifesi, F., Ajayi, O., Sileshi, G.W., Mowo, J.G., Kalinganire, A., Larwanou, M., Bayala, J., 2010. Evergreen Agriculture: a robust approach to sustainable food security in Africa. *Food Secur.* 2, 197–214.
- Giller, K.E., Cadisch, G., 1995. Future benefits from biological nitrogen fixation: an ecological approach to agriculture. *Plant Soil* 174, 255–277.
- Giller, K.E., Witter, E., Corbeels, M., Titttonell, P., 2009. Conservation agriculture and smallholder farming in Africa. The heretics' view. *Field Crops Res.* 114, 23–34.
- Govaerts, B., Verhulst, N., Castellanos-Navarrete, A., Sayre, K.D., Dixon, J., Dendooven, L., 2009. Conservation agriculture and soil carbon sequestration: between myth and farmer reality. *Crit. Rev. Plant Sci.* 28, 97–122.

- Gowing, J.W., Palmer, M., 2008. Sustainable agricultural development in sub-Saharan Africa: the case for a paradigm shift in land husbandry. *Soil Use Manage.* 24, 92–99.
- Gupta, R., Sayre, K., 2007. Conservation agriculture in South Asia. *J. Agric. Sci.* 145, 207–214.
- Guto, S.N., Pypers, P., de Ridder, N., Vanlauwe, B., Giller, K.E., 2011. Socio-ecological niches for minimum tillage and crop-residue retention in continuous maize cropping systems in smallholder farms of Central Kenya. *Agric. J.* 103, 1–11.
- Harris, F., 2002. Management of manure in farming systems in semi-arid West Africa. *Exp. Agric.* 38, 131–148.
- Hauser, S., Nolte, C., Carsky, R.J., 2006. What role can planted fallows play in the humid and sub-humid zone of West and Central Africa? *Nutr. Cycl. Agroecosyst.* 76, 297–318.
- Jama, B., Buresh, R.J., Place, F.M., 1998. Sesbania tree fallows on phosphorus-deficient sites: maize yield and financial benefit. *Agron. J.* 90, 717–726.
- Kang, B.T., Wilson, G.F., Siphens, L., 1981. Alley cropping maize (*Zea mays* L.) and *Leucaena leucocephala* Lam) in Southern Nigeria. *Plant Soil* 63, 165–179.
- Kassam, A., Friedrich, T., Shaxson, F., Pretty, J., 2009. The spread of Conservation Agriculture: justification, sustainability and uptake. *Int. J. Agric. Sustain.* 7, 292–320.
- Knowler, D., Bradshaw, B., 2007. Farmers' adoption of conservation agriculture: a review and synthesis of recent research. *Food Policy* 32, 25–48.
- Landers, J., 1999. How and why the Brazilian zero tillage explosion occurred? In: Stott, D.E., Mohtar, R.H., Steinhardt, G.C. (Eds.), *Sustaining the Global Farm. Selected Papers from the 10th International Soil Conservation Organization Meeting held May 24–29, 1999 at Purdue University and the USDA-ARS National Soil Erosion Research Laboratory.*, pp. 29–39.
- Luo, Z., Wang, E., Sun, O.J., 2010. Can no-tillage stimulate carbon sequestration in agricultural soils? A meta-analysis of paired experiments. *Agric. Ecosyst. Environ.* 139, 224–231.
- Mupangwa, W., Twomlow, S., Walker, S., 2012. Reduced tillage, mulching and rotational effects on maize (*Zea mays* L.) cowpea (*Vigna unguiculata* L. (Walp)) and sorghum (*Sorghum bicolor* L. (Moench)) yields under semi-arid conditions. *Field Crops Res.* 132, 139–148.
- Mwangi, W.M., 1997. Low use of fertilizers and low productivity in sub-Saharan Africa. *Nutr. Cycl. Agroecosyst.* 47, 135–147.
- Naudin, K., Scopel, E., Andriamandroso, A.L.H., Rakotosolofo, M., Andriamarosoa Ratsimbazafy, N.R.S., Rakotozandriny, J.N., Salgado, P., Giller, K.E., 2011. Trade-offs between biomass use and soil cover. The case of rice-based cropping systems in the Lake Alaotra region of Madagascar. *Exp. Agric.* 48, 194–209.
- Roose, E., Barthes, B., 2001. Organic matter management for soil conservation and productivity restoration in Africa: a contribution from Francophone research. *Nutr. Cycl. Agroecosyst.* 61, 159–170.
- Rusinamhodzi, L., Corbeels, M., van Wijk, M., Rufino, M.C., Nyamangara, J., Giller, K.E., 2011. Long-term effects of conservation agriculture practices on maize yields under rain-fed conditions: lessons for southern Africa. *Agron. Sustain. Dev.* 31, 657–673.
- Rusinamhodzi, L., Corbeels, M., Zingore, S., Nyamangara, J., Giller, K.E., 2013. Pushing the envelope? Maize production intensification and the role of cattle manure in recovery of degraded soils in smallholder farming areas of Zimbabwe. *Field Crops Res.* 147, 40–53.
- Sanchez, P.A., Buresh, R.J., Leakey, R.R., 1997. Trees, soils, and food security. *Phil. Trans. R. Soc. B* 352, 949–961.
- Sidibé, A., 2005. Farm-level adoption of soil and water conservation techniques in northern Burkina Faso. *Agric. Water Manage.* 71, 211–224.
- The Montpellier Panel, 2013. *Sustainable Intensification: A New Paradigm for African Agriculture.* Imperial College London, London.
- Thierfelder, C., Mombeyarara, T., Mango, N., Rusinamhodzi, L., 2013. Integration of conservation agriculture in smallholder farming systems of southern Africa. *Int. J. Agric. Sustain.* (in press).
- Tittonell, P., Giller, K.E., 2013. When yield gaps are poverty traps: the paradigm of ecological intensification in African smallholder agriculture. *Field Crops Res.* 143, 76–90.
- Tittonell, P., Vanlauwe, B., Leffelaar, P.A., Shepherd, K.D., Giller, K.E., 2005. Exploring diversity in soil fertility management of smallholder farms in western Kenya. II. Within farm variability in resource allocation, nutrient flows and soil fertility status. *Agric. Ecosyst. Environ.* 110, 166–184.
- Tittonell, P., Muriuki, A., Shepherd, K.D., Mugendi, D., Kaizzi, K.C., Okeyo, J., Verchot, L., Coe, R., Vanlauwe, B., 2010. The diversity of rural livelihoods and their influence on soil fertility in agricultural systems of East Africa – a typology of smallholder farms. *Agric. Syst.* 103, 83–97.
- Twomlow, S.J., Urolov, J.C., Jenrich, M., Oldrieve, B., 2008. Lessons from the field – Zimbabwe's Conservation Agriculture Task Force. *J. SAT Agric. Res.* 6, 1–11.
- Valbuena, D., Erenstein, O., Homann-Kee Tui, S., Abdoulaye, T., Claessens, L., Duncan, A.J., Gérard, B., Rufino, M.C., Teufel, N., Van Rooyen, A., Van Wijk, M.T., 2012. Conservation Agriculture in mixed crop–livestock systems: scoping crop residue trade-offs in sub-Saharan Africa and South Asia. *Field Crops Res.* 132, 175–184.
- Vanlauwe, B., Tittonell, P., Mukalama, J., 2006. Within-farm soil fertility gradients affect response of maize to fertilizer application in western Kenya. *Nutr. Cycl. Agroecosyst.* 76, 171–182.
- Vanlauwe, B., Bationo, A., Chianu, J., Giller, K.E., Merckx, R., Mkwunye, U., Ohiokpehai, O., Pypers, P., Tabo, R., Shepherd, K., Smaling, E., Woormer, P.L., Sanginga, N., 2010. Integrated soil fertility management: operational definition and consequences for implementation and dissemination. *Outlook Agric.* 39, 17–24.
- Vanlauwe, B., Kihara, J., Chivenge, P., Pypers, P., Coe, R., Six, J., 2011. Agronomic use efficiency of N fertilizer in maize-based systems in sub-Saharan Africa within the context of Integrated Soil Fertility Management. *Plant Soil* 339, 35–50.
- Verhulst, N., Sayre, K.D., Vargas, M., Crossa, J., Deckers, J., Raes, D., Govaerts, B., 2011. Wheat yield and tillage–straw management system × year interaction explained by climatic co-variables for an irrigated bed planting system in northwestern Mexico. *Field Crops Res.* 124, 347–356.
- Zingore, S., Johnston, A., 2013. The 4R Nutrient Stewardship in the Context of Smallholder Agriculture in Africa. In: Vanlauwe, B., Blomme, G., van Asten, P. (Eds.), *Agro-ecological Intensification of Farming Systems in the East and Central African Highlands.* Earthscan, UK (in press).